

Implementation and clinical utility of a Computer Aided Risk score for Mortality (CARM): A qualitative study

Judith Dyson*, Centre for Social Care, Health and Related Research, Birmingham City University, Ravensbury Building, Westbourne Road, B15 3TN, Judith.Dyson@bcu.ac.uk

Carolyn McCrorie, Yorkshire & Humber Patient Safety Translational Research Centre, Bradford Institute for Health Research

Jonathan Benn, Yorkshire & Humber Patient Safety Translational Research Centre, Bradford Institute for Health Research, School of Psychology, University of Leeds

Donald Richardson, York and Scarborough Teaching Hospitals NHS Foundation Trust,

Claire Marsh, Bradford Institute for Health Research

Gill Bowskill, Service User and Carer Research Group, Faculty of Health Studies, University of Bradford

Keith Double, Service User and Carer Research Group, Faculty of Health Studies, University of Bradford

Jean Gallagher, Service User and Carer Research Group, Faculty of Health Studies, University of Bradford

Muhammad Faisal, Faculty of Health Studies, University of Bradford, Yorkshire & Humber Patient Safety Translational Research Centre, Wolfson Centre for Applied Health Research, Bradford, UK

Mohammed A Mohammed, Faculty of Health Studies, University of Bradford, NHS Midlands and Lancashire Commissioning Support Unit, The Strategy Unit, Kingston House, West Bromwich, UK

Key words: Risk Score, Implementation, Mortality

Word count: 4045

Implementation and clinical utility of a Computer Aided Risk score for Mortality (CARM): A qualitative study

Abstract

Objectives: The Computer-Aided Risk score for Mortality (CARM) estimates the risk of in-hospital mortality following acute admission to hospital by automatically amalgamating physiological measures, blood tests, gender, age and Covid-19 status. Our aim was to implement the score with a small group of practitioners and understand their first-hand experience of interacting with the score in-situ.

Design: Pilot implementation evaluation study involving qualitative interviews.

Setting: This study was conducted in one of the two National Health Service hospital trusts in the North of England in which the score was developed.

Participants: Medical, older person and ICU/Anaesthetic consultants and specialist grade registrars (n=116) and critical outreach nurses (n=7) were given access to CARM. Nine interviews were conducted in total, with eight doctors and one Critical Care Outreach Nurse.

Interventions: Participants were given access to the CARM score, visible after login to the patients' electronic record, along with information about the development and intended use of the score.

Results: Four themes and fourteen subthemes emerged from reflexive thematic analysis; 1 current use (including support or challenge clinical judgement and decision making, communicating risk of mortality and professional curiosity), 2 barriers and facilitators to use (including litigation, resource needs, perception of the evidence base, strengths and limitations), 3 implementation support needs (including roll out and integration, access, training and education) and 4 recommendations for development (including presentation and functionality and potential additional data). Barriers and facilitators to use, and recommendations for development featured highly across most interviews.

Conclusions: Our in-situ evaluation of the pilot implementation of CARM demonstrated its scope in supporting clinical decision making and communicating risk of mortality between clinical colleagues and with service users. It suggested to us barriers to implementation of the score. Our findings may support those seeking to develop, implement or improve the adoption of risk scores.

Strengths and limitations

- Our development of CARM has benefited from the input of practitioners, service users and carers with regard to its content, presentation, use and implementation.
- Due to the scarcity of practitioner time during the period of Covid -19 our participant interviews were limited to n=9

- The CARM has been accessible to physicians on all medical wards. To date only nurses working in critical outreach have had access to the score.

INTRODUCTION

Unplanned or emergency medical admissions to hospital are common and involve patients with a broad-spectrum of disease and illness severity^{1,2}. The appropriate early assessment and management of such admissions can be a critical factor in ensuring safe and high-quality care. A number of risk scoring systems have been developed which may support the clinical decision-making process but few have been externally validated, implemented into routine practice and evaluated in situ³. We have developed a computer-aided risk of in-hospital mortality (CARM) score following emergency medical admission that automatically combines routinely collected, electronically recorded, clinical data (described elsewhere²) to support recognition of deterioration and therefore the prevent deaths attributable to poor clinical monitoring. The AUC C-statistic for CARM was 0.86 indicating good discrimination; it has been demonstrated to be more accurate than other similar risk scores³ and compares well with senior medical clinical judgement in identifying medical patients who were discharged alive or died in hospital (survival to discharge)⁴. CARM amalgamates the National Early Warning Score (NEWS2)⁵ with routinely taken blood test results, age, gender and more recently Covid-19 status (to make it relevant during the pandemic).

Uniquely, concurrent to the statistical development of CARM, healthcare practitioners, service users and carers contributed to the development of the score to establish views on the potential value, unintended consequences, concerns and implementation needs of the score⁶. Health technologies, such as risk scores, may falter due to poor implementation practices, poorly designed usability features and other contextual factors⁶. Therefore, i) as a preliminary to broader implementation of CARM in other hospitals sites, ii) to understand how potential users might interact with CARM in a clinical service environment and iii) to understand the contextual factors contributing to successful implementation and uptake, CARM was piloted at York and Scarborough Teaching Hospitals NHS Foundation Trust (YSTHNNFT) by integrating it into the electronic patient health record and making it available (on login) to authorised users.

Objective

Conduct a rapid in-situ qualitative evaluation to understand the first-hand experience of practitioners interacting with CARM and making clinical decisions relating to individual patients in clinical practice.

METHODS

Study Design

Presentation of the score: CARM is expressed as a two decimal point number from 0 (representing 0%) the lowest to 1 (representing 100%) being the highest risk of mortality. The score sat on the “home” screen of each patient record and by clicking on the score the user would be taken to a screen showing each component of CARM (that is, age, gender, NEWS elements and individual blood results). This is illustrated in figure 1 for an anonymised patient. The time the score was generated, NEWS2 score and components are shown on the left and the component blood tests are shown on the right. If users click the “chart” button they are taken to a trend chart of all CARM scores for the patient’s episode of admission.

Implementation of CARM at the Pilot Site: The practitioner group described in table 1, were informed of their access to CARM by way of an email circulated by the Chief Clinical Information Officer/Deputy Medical Director and clinical lead for the development of CARM (DR) on 20th November 2020. They were informed CARM would be live on 1st December 2020 and given links to publicly available, accessible video and online written information about CARM ([CARSs Research – Helping clinical staff make informed decisions about risk](#))⁷. This information describes the development, purpose and use of CARM. Email instructions included: i) CARM should not be used in isolation from other sources of information, ii) CARM was intended to augment (not replace) clinical decision-making and iii) CARM relies upon processing existing information to improve situational awareness, rather than introducing new sources of information.

Interview questions: Our interview schedule was informed by the study aim, to understand the first-hand experience of practitioners exposed to CARM in their daily practice and clinical decision-making. Questions included general experience and perceptions of impact, value, problems and presentation.

Setting

This study took place at YSTHNHSFT, a two-site 1200-bed academic acute care trust in the UK and one of the trusts in which CARM was developed. The trust has an in-house electronic patient record development team and a mature infrastructure tried and tested to combine NEWS2 with blood test results.

Participants

CARM was activated for physicians graded specialist registrar and above working in medical environments and for ICU consultants and nurses on the critical care outreach team (CCOT) all of

whom may be called to assess and potentially admit deteriorating patients to ICU. This group comprised the potential participants of the study and the numbers of each group with access to CARM are indicated in table 1. Our aim was to sample perspectives and experiences drawn from a range of different user groups and we expected approximately 10-15 participants would be sufficient to achieve data saturation ⁸.

Table 1: Characteristics of those with access to CARM

Practitioner group	Number exposed to CARM
Medical Consultants (C)	50
Elderly Medicine Consultants	15
Medical Specialist Registrars	30
Elderly Medicine Specialist Registrars	7
Consultant Intensivists on CCOT	14
CCOT Nurses	7
Total	123

Patient and public involvement


We have sought to co-design CARM with service users and carers as part of the project team as well as participants^{6,9}. In the study reported here, we had the input of three team members (GB, KD, JG) from the “Service User and Carer involvement in Research Group” at the University of Bradford, who have been involved in the development of CARM over the last four years. Our service user team members were involved in interpretation of data from the patient perspective, reading and commenting on results and contributing ideas to include in the discussion and writing of this paper. We presented our data to the clinical governance group of the hospital to check our results made sense from a clinical and organisational perspective.

Procedure

Once participants had been exposed to CARM for four months, a further email was sent by DR inviting them to participate in an interview and share their experiences of CARM. A participant information sheet about the aim of the study was attached with instructions to respond to JD or CMc. Interviews took place on the telephone (due to Covid-19 social distancing guidelines) and were audio recorded and transcribed verbatim. Two subsequent reminders were sent. All volunteers were interviewed by either JD or CMc, both experienced, post-doctoral qualitative health service researchers, neither of whom had any professional or personal knowledge or contact with the hospital staff included. Volunteers came from all three clinical groups (anaesthetist/intensivist, medical and nursing) leaving subsequent purposive sampling unnecessary. Prior to interviews

questions were answered and verbal consent taken. Interviews were held between May and July 2021.

Figure 1: Presentation of CARM for Practitioners



Analysis

Interviews were professionally transcribed and analysed within NVivo (v12). Qualitative data analysis proceeded in line with Braun and Clarke's six step process (2006), drawing upon subsequent formulations of Reflexive Thematic Analysis (RTA) ¹⁰⁻¹². Reflexive Thematic Analysis was selected as an approach well-suited to flexible exploration of user's perceptions and experiences, whilst balancing subjectivity, the analytic process and the data itself through iterative reflection¹¹. Data analysis proceeded in parallel with data collection until saturation was achieved (no further novel initial themes emerged). Once preliminary interviews had been completed, the research group first familiarised themselves with the transcribed text. Initial inductive coding was then undertaken by CMc and JD, initially independently, in order to generate candidate themes. Themes were identified at the semantic level initially, with subsequent refinement focusing on latent concepts. Following reflection on the independent coding, a convergent coding framework and commentary was produced by JD, integrating the prior codes which was subsequently refined in preparation of the report by CMc, JB and JD.

Ethics

The protocol for the study was reviewed by Leeds West NRES Committee in October 2017 (17/YH/0367). In accordance with regional guidelines, the research ethics committee deemed the study was a service evaluation and formal ethical review was not required. The service evaluation was registered with York Teaching Hospital NHS Foundation Trust in January 2020.

RESULTS

Nine interviews were conducted and took between 16 and 45 minutes (mean duration 31 minutes).

Characteristics of the sample

There were nine participants, eight doctors and one CCOT nurse. A summary of participant characteristics is presented in Table 2. Due to the small sample drawn from a single hospital we have not linked these details to participant ID (presented in frequency table and in subsequent quotations) to preserve anonymity. Rather, we have ordered participants randomly in table 2. General medical participants specified specialities included renal, emergency, gastroenterology and respiratory medicine.

Table 2: Participant characteristics

Role (experience in years)	Specialty	Exposure to/degree of adoption of CARM
Consultant (13)	General Medicine	Views the score to confirm/challenge clinical judgement
Medical Consultant (4)	General medicine	Check the score after making clinical judgement
Consultant (23)	ICU/ anaesthetics	Views the score out of professional curiosity
Medical Consultant (18)	ICU/ anaesthetics	Views CARM to confirm/challenge clinical judgement when called to assess deteriorating patients on medical ward
Senior Nurse (1.5)	Critical Outreach team	Views the score to confirm/challenge clinical opinion when assessing deteriorating patients on medical ward
Medical Consultant (12)	ICU/ anaesthetics	Views CARM when called to assess deteriorating patients on medical ward
Medical Consultant (10)	General medicine	Views the score out of professional curiosity and to confirm clinical judgement
Specialist Registrar (3)	General Medicine	Views the score to confirm/challenge clinical judgement
Medical Consultant (14)	General medicine	Views the score to confirm/challenge clinical judgement

Findings

There were four themes and fourteen subthemes; theme 1 current use (including support or challenge clinical judgement and decision making, communicating risk of mortality and professional curiosity), theme 2 barriers and facilitators to use (including litigation, resource needs, perception of the evidence base and strengths and limitations), theme 3 implementation support needs (including roll out and integration, access and training and education) and theme 4 recommendations for development (including presentation and functionality and potential additional data). These are illustrated in figure 2 and presented in turn below with any differences between groups described within the narrative. The frequency with which participants spoke about any one theme is presented in table 3. Barriers and facilitators to use and recommendations for development featured highly across most interviews.

Figure 2: Thematic map of practitioner views on the clinical utility and implementation support needs of CARM




Table 3: Frequency with which themes were discussed according to participant

Participant	Current Use	Barriers and facilitators to use	Implementation support needs	Recommendations for development
1	6	7	2	7
2	1	6	7	8
3	4	6	2	3
4	7	8	7	9
5	6	6	3	2
6	4	8	0	5
7	1	2	2	0
8	8	9	3	5
9	4	5	3	3

1. Current use

This theme described how those practitioners currently exposed to CARM reported using it.

i) To support or challenge clinical judgement and decision making: Most participants explained how CARM confirmed (reassured) or prompted them to reconsider their clinical judgement and care decisions. There were two key areas where CARM supported judgement: ceiling of care decisions and escalation (e.g., admission to ICU). Some reported CARM prompted earlier decisions than they might have made otherwise.

“End of life, or people who are very sick. If you’ve got a, if you’ve got a CARM that’s sort of, that supports your clinical judgement and assessment, then that’s, there’s a degree of reassurance in there, so for me that is a benefit.” P5

“In that lady it prompted a Do Not Resuscitation discussion with her family members. I’d say the change would be that I’m making those decisions earlier.” P7

ii) As a means of communicating risk of mortality: CARM was often used as a means of communication with colleagues to support optimal patient care and for teaching purposes. However, for patients and relatives the quantitative nature of chance of survival was considered particularly useful.

“Because saying to someone, “oh, you’ve got a very slim chance of surviving,” initial reaction is often. . . “Well, I want everything . . . do what you can.” But then you explain well, actually, it’s less than 10% even if we do everything . . . “Oh, right, OK, I understand that now.” It sort of makes things a bit easier.” P8

iii) Out of professional curiosity: Some practitioners didn’t include CARM in their clinical decision making, they simply looked out of interest and considered how it *might* fit into their everyday practice. The reasons given were they saw the score as still being in development as part of a research project, the score had not yet been embedded into clinical systems, because only few practitioners had access to the score and because they did not have enough background information on the score to know how to use it.

“I’m treating it as a research tool . . . I’m mindful that I can’t give it as much weight . . . I know it’s been validated, but in my mind it’s still a research tool, so therefore it’s not going to heavily, heavily play, necessarily, on my decisions.” P3

2. Barriers and facilitators to use

There were a number of elements relating to CARM users saw as either a barrier or a facilitator.

i) Litigation was a concern for a small number of participants; they believed acknowledging without taking action on CARM might leave them open to criticism. More often, criticism for failure to act was seen as an appropriate response; if a score indicates a risk action should be taken.

“If you write something in the notes about this and then you choose to acknowledge but not do anything about it, it’s, yeah, you can be hauled over the coals for it, couldn’t you, in a court.” P4

ii) Resource needs: Some participants expressed the possibility of CARM resulting in more patients needing intensive care and the potential for the need to clinically prioritise the use of those beds. Concerns about resources were also often mitigated with views that if deteriorating patients were identified sooner (using CARM) this may result in more efficient care (e.g., earlier but shorter stays in the ICU). One participant thought if CARM identified deterioration early enough it would result in more referrals to the critical outreach team, and this could *prevent* an admission to ICU:

“If we need the resources . . . shout about it and say we need more ICU beds . . . [CARM may] lead to more referrals direct to the critical care team . . . the earlier we see them the more likely it is that we can prevent the deterioration to ever need critical care.” P1

iii) Perception of the evidence base for CARM: Some participants were unfamiliar with the evidence underpinning the CARM and questioned the validity of the score and were therefore reluctant to adopt the into their clinical decision making CARM.

“If I was, if I was looking at a score like this, I’d want to know . . . how it’s been validated, what groups it’s been validated in.” P8

One participant was concerned CARM may over-estimate mortality.

“I think overestimating mortality risk, I think I have come across situations where I’ve thought that maybe it was doing that.” P1

When practitioners believed the score under-estimated the risk of mortality it was when the score was considered in isolation. Most practitioners understood CARM was only one part of the clinical picture, one tool among many to support clinical decision making.

“You’d need to see the whole, you know, the overall condition and trend of trajectory of the patient’s condition was.” P3

iv) Strengths and limitations of CARM were expressed, often relative to NEWS. Participants considered CARM more accurate than NEWS but thought the less frequent calculation and presentation a limitation.

“The CARM takes into, takes into account, not only just the observations but . . . the demographic, the age of a patient, and also the . . . biochemistry in there it’s not just as, as two-dimensional as the observation chart . . . [CARM] is able to give a more rounded overview of how mortality is predicted, not just a simple observational chart.” P7

“It’s not quite as dynamic a score as the NEWS score because . . . we’re not measuring the various blood parameters every day.” P3

3. Implementation support needs

This theme related to the actions needed to be taken to fully implement and embed CARM into routine daily practice.

i) Roll out and integration: One participant considered roll out should be part of a large-scale research project:

“The team goes for a big research grant . . . all-singing all-dancing off we go with a fanfare . . . that’s what I’d prefer.” P1

Other suggestions included small group discussions and time to become familiar with the score, embedding it into practice, systems and processes.

“A few of us getting together and talking about it, or [Clinical lead] or whoever . . . coming to work through some with you in the hospital would be the most useful. Just a gradual rollout” P2

“We have to get a feel for it . . . I think it does take a little bit of time to just sort of embed that into your, into your practice.” P7

Participants were not sure what actions to take according to the score. Some participants suggested this might be dealt with through some sort of escalation policy or protocol.

“What are you doing with this information?” P5

“Currently . . . it’s not part of the escalation policy.” P4

ii) Access: Some participants offered opinions on who might best access the system. Some thought anyone involved in taking and responding to physiological observations, others said due to the nature of the score (a predictor of mortality) it should be restricted to medical staff.

“Because it’s to do with prognosis, I’d probably, without sounding too patronising, I’d probably keep it to medical staff.” P8

“The healthcare assistants do a lot of the observations, so it’s really important that they’re empowered to speak up about these things.” P2

Although participants generally appreciated how the score was displayed, logging onto the system in different areas could be arduous.

"I don't always get access to the CARM score because, because, mainly because of the IT side of things. So, I've got to go and login on my login . . . if I'm going up to a ward, if I'm going to a computer that's that I've not been to before, you might spend ten plus minutes logging, you know, for the computer to learn your permissions." P3

iii) **Training and education**

Most participants suggested training and education to support the implementation of CARM. Content included the background to the score (including validation), how the score is calculated (the components), how it is different to the NEWS and what actions to take according to the score.

"I'd like to know more about it really." P9

"The background to it and how it's calculated at and why it gives extra weight for a NEWS score [and] what do you do with that information?" P4

4. Recommendations for development

Participants had a number of suggestions for the development of CARM.

i) Presentation: Some suggested having access to the CARM in different formats. Participants linked with ICU suggested it would be a good idea for the critical outreach team to see the CARM for all patients in the hospital in order to prioritise the patients they might visit.

"So currently . . . we can filter high NEWS scores across the Trust . . . it would probably be quite useful . . . see who's got a high-risk score and then be able to go and be a bit more proactive." P4

Several participants suggested CARM being accessible electronically via mobile telephones, tablets and other devices and in hard copy in the ward environment. Some suggested a visual "flag" if CARM suggested the need for a review or a traffic light system to prioritise response.

". . . could be on mobile 'phones . . . on the ward whiteboards" P1

"The traffic light system . . . it might actually make, draw your attention a little bit more, rather than a number on the bottom of the screen." P2

Participants particularly appreciated the option to see previous CARM scores presented as a trend graph:

"Being able to look at changes over time, because to me that's one of the most useful things" P2

Some participants suggested additional functionality to identify which of the various elements contributing to CARM (e.g., which physiological element or blood test) had led to the score change.

Others wanted an indication of when the score was refreshed with new data.

“So, you do get patients whose physiological parameters are abnormal when their blood tests are normal. Or those blood tests are outdated . . . you just would have to be very clear about what the timescale was.” P6

ii) Potential additional data to include in CARM: Whilst our focus was implementation and perceived utility, there were suggestions about additional data to include in CARM. We include this as a theme, not only because they may be relevant to the future development of CARM, but because offering practitioners the rationale for not including these elements may be important to wider scale implementation. Diagnosis was frequently suggested, particularly if this was life limiting. Other suggestions were the addition of comorbidities, frailty data, nursing concerns and previous hospital admissions, all of which participants considered might aid mortality prediction.

“Comorbidities . . . single organ failure, multi-organ failure or heart failure, liver failure . . . frailty.” P8

“Nursing concern . . . there are some Trusts that would add an extra 3 points [to NEWS] for nursing concern.” P4

“Frequency of hospitalisation.” P5

DISCUSSION

Our approach to the development of CARM has involved a process of co-design with healthcare professionals and service users⁶. Previous papers discuss the development and validation² and the potential value, unintended consequences, concerns and predicted implementation needs⁶ of the CARM. Building on these findings, in this study we report the small pilot implementation and evaluation of CARM based upon practitioners’ experiences of accessing it in clinical practice. The majority of the group interviewed had actively used CARM to support or challenge their decision making and to communicate about a patient’s status. Practitioners identified factors that helped or hindered their engagement with CARM along with their implementation support needs and recommendations for additional functionality of the score. Finally, practitioners suggested additional clinical data for consideration for inclusion in CARM, the potential for which we discuss below.

Our findings relating to current use of CARM, barriers and facilitators and implementation support needs feed into the next steps for implementing CARM across the current hospital environment, prior to scaling up. In developing CARM our aim was to identify and prevent deaths attributable to poor clinical monitoring. Our work here (and previously⁶) suggests CARM goes beyond this and supports clinical decisions relating to both escalation and ceiling of care. CARM aids communication with colleagues and service users. We identified barriers to CARM and implementation support

needs we can seek to address prior to the next stage. Barriers expressed indicate poor engagement or inadequacy of our existing implementation materials. In line with previous research relating to risk scores on computerised systems¹³ our participants had difficulty viewing contributing elements and had suggestions to enhance the display. Our findings illustrate the need for careful implementation of risk scores, early consideration and design of an implementation package and the need to understand the requirements for support and training.

Some of our findings relating to implementation concur with other seeking to implement risk scores. For example, there is a modest literature exploring barriers to NEWS adoption summarised in two recent reviews and including lack of training^{14 15} (identified as a need by our participants) poorly designed documentation systems¹⁵ (which we sought to address through co-design). Other barriers identified by others included high workload^{14 15} and lack of skills¹⁵. Smith et al¹³ investigated barriers to taking and recording physiological measures using an electronic system (most such studies involve paper-based systems). Barriers specific to the electronic system included difficulty viewing elements contributing to an elevated score and a poor display format¹³. Similarly, our participants suggested a bolder means of presenting the score (e.g., traffic lights) and clearer links to the elements contributing CARM. Such barriers exist with non-mortality related computer risk scores and in other environments, for example implementation needs relating to an infection risk score in care homes were similar to ours, including, training and education needs and challenges in integrating the score into existing systems¹⁶. This suggests that findings reported here may be transferrable to the implementation of other risk scores.

There were strengths and limitations to this study. To the best of our knowledge, CARM is the only published risk score to have involved service providers and users (as both research partners and participants) in its development as well as implementation. We did all possible to follow good practice guidelines for co-working including working as equals and meaningful, early engagement¹⁷⁻²⁰. We have received feedback from our service user colleagues²¹ and we report here and elsewhere⁶ on the impact of practice and service user views on the score. Because of the clear value that involvement has added to both the development of CARM and in this study, we will consider formal evaluation of co-design in follow-up projects where involvement will continue to be the bedrock to our work. A population of only 123 potential participants (those with access to CARM) all of whom are busy clinicians, during a global pandemic made recruitment challenging and we ceased recruitment after interviewing nine participants. The implementation of complex interventions requires preliminary evaluation of the intervention within the context in which it is implemented and how it contributes to support decision making in the real world²². We sought to achieve this through flexible approaches to the timings of interviews and acknowledgement of the potential

need to reschedule due to clinical shift changes or pressures. We met the balance of gaining meaningful data whilst avoiding taking more time than necessary from potential participants by concurrent data analysis and collection and vigilance in abiding to the guiding principles of data saturation²³. The only nurses exposed to CARM were critical outreach nurses; we did not include nurses working in other medical settings. Our findings suggest CARM has use for nurses and our future work will include this group, however, the transferability of our findings is limited to selected groups. Finally, although we captured self-reported exposure/degree of adoption with CARM (table 2) and we appear to have a range of responses, there may have been an element of selection bias whereby those volunteering for interviews were those the most engaged with CARM.

CONCLUSION

Our in-situ evaluation of the pilot implementation of CARM has demonstrated its scope in supporting clinical decision making and its use in communicating risk of mortality between clinical colleagues and with service users. We have gained insights into barriers to full implementation and adoption, implementation needs and suggestions for the development of the score in terms of presentation, functionality and content. Our findings may support those seeking to develop, implement or improve compliance with other risk scores (including NEWS).

Contributorship statement: MM and DR had the original idea for this work. JD, CMc, JB and DR were the leads for this implementation/utility study. CMa led the service user contribution. JD, CMc, JB and MM wrote the first draft of this paper and all authors (including GB, KD, JG, DR and MF) assisted in redrafting and have approved the final version.

Competing Interests: None declared.

Funding statement: This research is funded by the National Institute for Health Research (NIHR) Yorkshire and Humber Patient Safety Translational Research Centre (NIHR Yorkshire and Humber PSTRC). The views expressed in this article are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

Data sharing statement: Anonymised qualitative data are available from the corresponding author.

References

1. Downey C, Tahir W, Randell R, et al. Strengths and limitations of early warning scores: a systematic review and narrative synthesis. *International Journal of Nursing Studies* 2017;76:106-19.
2. Faisal M, Scally AJ, Jackson N, et al. Development and validation of a novel computer-aided score to predict the risk of in-hospital mortality for acutely ill medical admissions in two acute hospitals using their first electronically recorded blood test results and vital signs: a cross-sectional study. *BMJ open* 2018;8(12):e022939.
3. Brabrand M, Folkestad L, Clausen NG, et al. Risk scoring systems for adults admitted to the emergency department: a systematic review. *Scandinavian journal of trauma, resuscitation and emergency medicine* 2010;18(1):1-8.
4. Faisal M, Khatoun B, Scally A, et al. A prospective study of consecutive emergency medical admissions to compare a novel automated computer-aided mortality risk score and clinical judgement of patient mortality risk. *BMJ open* 2019;9(6):e027741.
5. Royal College of Physicians. National Early Warning Score (NEWS): Standardising the assessment of acute illness severity in the NHS - Report of a working Party, 2012.
6. Dyson J, Marsh C, Jackson N, et al. Understanding and applying practitioner and patient views on the implementation of a novel automated Computer-Aided Risk Score (CARS) predicting the risk of death following emergency medical admission to hospital: qualitative study. *BMJ open* 2019;9(4):e026591.
7. Richardson D. Helping clinical staff make informed decisions about risk [Available from: <https://carsresearch.org/2022>].
8. Guest G, Bunce A, Johnson L. How many interviews are enough? An experiment with data saturation and variability. *Field methods* 2006;18(1):59-82.
9. Faisal M, Richardson D, Scally A, et al. Performance of externally validated enhanced computer-aided versions of the national early warning score in predicting mortality following an emergency admission to hospital in England: A cross-sectional study. *BMJ open* 2019;9(11):e031596.
10. Braun V, Clarke V. Thematic analysis In Cooper H, Camic PM, Long DL, Panter AT, Rindskopf D, & Sher KJ (Eds.), *APA handbook of research methods in psychology, Vol 2: Research designs: Quantitative, qualitative, neuropsychological, and biological* (pp. 57–71). *Washington, DC: American Psychological Association*[Google Scholar] 2012
11. Braun V, Clarke V. Reflecting on reflexive thematic analysis. *Qualitative Research in Sport, Exercise and Health* 2019;11(4):589-97.
12. Byrne D. A worked example of Braun and Clarke's approach to reflexive thematic analysis. *Quality & Quantity* 2021:1-22.
13. Smith D, Cartwright M, Dyson J, et al. Barriers and enablers of recognition and response to deteriorating patients in the acute hospital setting: A theory-driven interview study using the Theoretical Domains Framework. *Journal of Advanced Nursing* 2021;77(6):2831-44.
14. Olsen SL, Søreide E, Hillman K, et al. Succeeding with rapid response systems—a never-ending process: a systematic review of how health-care professionals perceive facilitators and barriers within the limbs of the RRS. *Resuscitation* 2019;144:75-90.
15. Wood C, Chaboyer W, Carr P. How do nurses use early warning scoring systems to detect and act on patient deterioration to ensure patient safety? A scoping review. *International journal of nursing studies* 2019;94:166-78.
16. Dowding D, Russell D, McDonald MV, et al. "A catalyst for action": Factors for implementing clinical risk prediction models of infection in home care settings. *Journal of the American Medical Informatics Association* 2021;28(2):334-41.

17. Greenhalgh T, Jackson C, Shaw S, et al. Achieving research impact through co-creation in community-based health services: literature review and case study. *The Milbank Quarterly* 2016;94(2):392-429.
18. Involve. Guidance on co-producing a research project. INVOLVE [Available from: <https://www.invo.org.uk/posttypepublication/guidance-on-co-producing-a-research-project/2021>].
19. Flinders M, Wood M, Cunningham M. The politics of co-production: risks, limits and pollution. *Evidence & Policy: A Journal of Research, Debate and Practice* 2016;12(2):261-79.
20. Shimmin C, Wittmeier KD, Lavoie JG, et al. Moving towards a more inclusive patient and public involvement in health research paradigm: the incorporation of a trauma-informed intersectional analysis. *BMC Health Services Research* 2017;17(1):1-10.
21. University of Bradford. Helping clinical staff make informed decisions about risk.
22. Skivington K, Matthews L, Simpson SA, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *bmj* 2021;374
23. Francis JJ, Johnston M, Robertson C, et al. What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychology and health* 2010;25(10):1229-45.