



Effects of the FIFA 11 + Program on Physical Fitness in Youth and Adult Soccer Players: A Systematic Review and Meta-analysis

Ibnu Noufal Kambitta Valappil¹ · Karuppasamy Govindasamy² · Gavoutamane Vasanthi¹ · Masilamani Elayaraja¹ · Cain C. T. Clark³ · Koulla Parpa⁴ · Borko Katanic⁵ · Hüseyin Şahin Uysal⁶ · Hassane Zouhal^{7,8} · Urs Granacher⁹ 

Accepted: 16 October 2025
© The Author(s) 2025

Abstract

Background Soccer is a high-intensity sport that requires high levels of physical fitness, including balance, change of direction (CoD), speed and power. The FIFA 11 + program has been widely promoted to enhance physical fitness and reduce injury occurrence.

Objective This meta-analysis set out to examine how the FIFA 11 + program, implemented as a warm-up versus conventional warm-up (soccer-specific and alternative warm-ups), impacts physical fitness attributes in youth and adult soccer players.

Methods After a priori defined inclusion and exclusion criteria, 17 intervention studies with 611 male and female soccer players (Tier 2–4) aged 9–29 years were eligible to be included. The FIFA 11 + program implemented in the warm-up was contrasted with conventional warm-up programs (control) on outcome measures such as dynamic balance, CoD speed, linear sprint and proxies of muscle power (vertical jump height). The influence of potential moderators (e.g., training duration, frequency, session duration, age, sex, training and performance calibre) on study outcome measures was examined using subgroup analyses with the median split method.

Results Findings demonstrated small-to-moderate improvements in favour of FIFA 11 + compared with conventional warm-ups on dynamic balance (small standardized mean differences [SMDs] = 0.37, $p < 0.001$, heterogeneity [I^2] = 7), CoD speed (moderate SMDs = -0.65, $p = 0.005$, $I^2 = 84$), and vertical jump height (small SMDs = 0.56, $p < 0.001$, $I^2 = 71$). Results from the sub-analyses showed that, for dynamic balance, shorter training durations (<9 weeks) produced larger effects than longer durations (≥ 9 weeks) (SMDs = 0.62 versus SMDs = 0.17). For vertical jump height, <9 weeks also yielded greater improvements (SMDs = 0.79 versus SMDs = 0.26). In terms of weekly training frequency, ≥ 3 sessions/week elicited larger gains in change-of-direction speed (SMDs = -1.05 versus SMDs = -0.12) and vertical jump height (SMDs = 0.73 versus SMDs = 0.01) compared with <3 sessions/week. Regarding participant characteristics, players aged ≥ 18 years showed greater improvements than those <18 years in change-of-direction speed (SMDs = -1.45 versus SMDs = -0.06) and vertical jump height (SMDs = 0.64 versus SMDs = 0.22). For sex differences, males experienced greater benefits than females in change-of-direction speed (SMDs = -0.79 versus SMDs = -0.04) and vertical jump height (SMDs = 0.54 versus SMDs = 0.09). Finally, higher-level players ($\geq$ tier 3) demonstrated greater improvements in vertical jump height than lower-tier players (< Tier 3) (SMDs = 0.75 versus SMDs = 0.01). The observed benefits were statistically significant but generally of small-to-moderate magnitude and affected

✉ Urs Granacher
Urs.Granacher@sport.uni-freiburg.de

¹ Department of Physical Education and Sports, Pondicherry University, Puducherry 605014, India

² Department of Sports, Recreation and Wellness, Symbiosis International (Deemed University), Hyderabad Campus, Modallaguda (V), Nandigama (M), Rangareddy, Telangana 509217, India

³ College of Life Sciences, Birmingham City University, Birmingham B15 3TN, UK

⁴ Faculty of Sport and Exercise Science, UCLan University of Cyprus, 7080 Pyla, Cyprus

⁵ Montenergin Sports Academy, Podgorica, Montenegro

⁶ Department of Physical Education and Sport, Faculty of Sport Sciences, Burdur Mehmet Akif Ersoy University, 15030 Burdur, Turkey

⁷ Institut International des Sciences du Sport (2I2S), 35850, Irodouer, France

⁸ Laboratoire optimisation de la performance sportive, centre national de la médecine et des sciences des sports, (LR09SEP01) Tunis, Tunisia


⁹ Department of Sport and Sport Science, Exercise and Human Movement Science, University of Freiburg, 79102 Freiburg, Germany

by study heterogeneity and program implementation differences. Most studies included male participants, limiting generalizability to female and underrepresented populations. Several studies also lacked rigorous methodological design, particularly in allocation, concealment and blinding. Reporting of training and demographic variables was often incomplete. These limitations highlight the necessity for rigorously designed, low-bias randomized controlled trials with standardized implementation of the FIFA 11 + program and thorough reporting to enhance the reliability of causal conclusions and improve clinical interpretation.

Conclusion The FIFA 11 + program was more effective than conventional soccer warm-up programs to enhance soccer players' physical fitness (i.e., dynamic balance, CoD speed, vertical jump height). A shorter training duration (<9 weeks) and more weekly training sessions (≥ 3 sessions/week) induced larger performance effects. FIFA 11 + was more effective in older (≥ 18 years) and male players and in players of higher performance calibre ($\geq$ Tier 3). These findings underscore the versatility of the FIFA 11 + program to improve soccer players' physical fitness, supporting its integration into the warm-up of regular soccer training sessions.

Protocol Registration <https://www.crd.york.ac.uk/PROSPERO/view/CRD42025633810>, PROSPERO: CRD42025633810.

Graphical Abstract



Key Points

The FIFA 11 + program has primarily been applied to prevent injuries in youth and adult soccer. Less is known about the performance-enhancing effects of FIFA 11 + when implemented as part of a warm-up.

This meta-analysis revealed that FIFA 11 + is effective in improving dynamic balance, linear sprint, change of direction speed and vertical jump height in youth and adult male and female soccer players aged 9–29 years.

FIFA 11 + seems to be particularly effective with a training duration of <9 weeks and ≥ 3 weekly training sessions. FIFA 11 + was more effective in older (≥ 18 years) and male players, and in players of higher performance calibre ($\geq$ Tier 3).

1 Introduction

Optimizing player preparation strategies remains a key focus in soccer, with growing interest in how structured warm-up routines may impact on performance enhancement and injury prevention [1, 2]. Evidence suggests that well-designed warm-up programs can improve neuromuscular control, enhance agility and reduce the incidence of non-contact injuries [3, 4]. Moreover, integrating sport-specific exercises within these routines may lead to acute performance benefits and long-term physiological adaptations [5–7]. In 2006, the World Soccer Association (FIFA) developed a training program called FIFA 11 +, which contains the multimodal components of neuromuscular training [8]. The FIFA 11 + program is particularly suited to be included as a warm-up program in soccer-specific training sessions in youth and adult male and female soccer players [9–12].

Originally developed as an injury prevention program, the FIFA 11 + program has also been effective in enhancing players' physical fitness [13]. It is important to note that these performance benefits may arise through two distinct pathways: (1) acute effects, such as neuromuscular facilitation and improved readiness when the program is applied immediately as a warm-up [5, 6, 14], and (2) chronic adaptations, including long-term improvements in measures of strength, balance and power when the program is implemented consistently over several weeks [4]. Considering the well-established chronic benefits of the FIFA 11 + program for injury prevention and performance enhancement, it has been adopted by different sports such as volleyball [15], basketball [16, 17], rugby [18, 19] and futsal [3, 20], becoming a cornerstone of pre-match preparation and training. The performance-enhancing effects of the FIFA 11 + are believed to be underpinned by neuromuscular adaptations, including improved motor unit recruitment, increased lower-limb and core muscle strength, enhanced proprioception and better intermuscular coordination [21–23]. These adaptations collectively contribute to superior balance, faster and more efficient change-of-direction movements, and increased jumping and sprinting capabilities [4, 24].

The most recent systematic review and meta-analysis examining the chronic effects of the FIFA 11 + on physical fitness, particularly change of direction (CoD) speed in soccer and futsal, was conducted in 2021 [24]. The analysis included 15 studies, and the authors concluded that the FIFA 11 + training program has the potential to significantly enhance CoD performance. However, the previous review incorporated five studies involving futsal players, whereas the present study focuses exclusively on soccer players. It is important to note that futsal differs substantially from soccer in terms of physiological demands, match-play characteristics, technical-tactical requirements and overall game dynamics, which may limit the generalizability of findings across the two sports [25–27]. Given that, 4 years have passed since the last meta-analysis on this topic, a

Table 1 Detailed search strategies from selected databases

Database	Search strategy
SCOPUS	('FIFA 11 + ' OR 'FIFA 11 plus' OR 'FIFA 11 + program*' OR '11 + ' OR 'FIFA 11 plus program*') AND ('football player*' OR 'soccer player*' OR 'football' OR 'soccer') AND ('speed' OR 'sprint' OR 'agility' OR 'change of direction' OR 'change-of-direction' OR 'COD' OR 'balance' OR 'jump' OR 'plyometric*' OR 'motor skill*' OR 'muscle power' OR 'athletic performance')
PUBMED	('FIFA 11 + ' OR 'FIFA 11 plus' OR 'FIFA 11 + program*' OR '11 + ' OR 'FIFA 11 plus program*') AND ('football player*' OR 'soccer player*' OR 'football' OR 'soccer') AND ('speed' OR 'sprint' OR 'agility' OR 'balance' OR 'jump' OR 'plyometric*' OR 'change of direction' OR 'change-of-direction' OR 'COD' OR 'motor skill*' OR 'muscle power' OR 'physical fitness' OR 'athletic performance'[MeSH])
WEB OF SCIENCE	TS=((('FIFA 11 + ' OR 'FIFA 11 plus' OR 'FIFA 11 + program*' OR '11 + ' OR 'FIFA 11 plus program*') AND ('football player*' OR 'soccer player*' OR 'football' OR 'soccer') AND ('speed' OR 'sprint' OR 'agility' OR 'change of direction' OR 'change-of-direction' OR 'COD' OR 'balance' OR 'jump' OR 'plyometric*' OR 'motor skill*' OR 'muscle power' OR 'physical fitness' OR 'athletic performance'))

Table 2 Selection criteria used in the meta-analysis

Category	Inclusion criteria	Exclusion criteria
Population	Male and female soccer players aged > 9+	Athletes from different sports other than soccer
Intervention	FIFA 11+ warm-up program	FIFA 11+ program applied as full training session and not as a warm-up; other training modalities than the FIFA 11+
Comparator	Active control group (soccer-specific and conventional warm-up)	Passive control groups
Outcomes	At least one measure of physical fitness including dynamic balance, linear sprint and CoD speed, agility, power (vertical jump height)	No physical fitness data, only injury prevention data
Study design	Randomized controlled trials (RCTs)	Cross-sectional studies, controlled studies, case studies and observational studies

re-evaluation of the scientific literature is warranted to provide a more comprehensive and up-to-date synthesis of the available evidence on FIFA 11+ effectiveness.

While the FIFA 11+ has shown overall positive effects on physical fitness [4], the magnitude of these benefits may depend on contextual factors such as training frequency, program duration, player age, sex and performance calibre [28, 29]. For example, higher training frequency could accelerate neuromuscular adaptations [23], older players may respond differently due to greater training experience and physical maturity, and the performance level of a player may impact on the capacity to translate training stimuli into performance

gains [30]. However, these moderating influences remain insufficiently explored in the existing literature. In this study, training and performance calibre was prioritized as a key moderator because players' competitive tier reflects not only their technical-tactical proficiency but also their physical conditioning and adaptability to structured training, making it a critical determinant of how effectively the FIFA 11+ program translates into performance outcomes [14, 31]. In contrast, variables such as session duration primarily reflect total program exposure rather than athletes' intrinsic adaptive capacity. For example, longer warm-up protocols may increase training load without yielding greater




Fig. 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram detailing identification, screening, eligibility criteria and included studies at each stage of the systematic review

performance improvements if baseline fitness and skill level remain unaccounted [14]. Focusing on performance calibre allows the analysis to capture how athletes' preparedness and adaptive capacity influence the effectiveness of the FIFA 11 + program. Addressing this gap through subgroup analyses can provide a deeper understanding of the conditions under which the FIFA 11 + is most effective, thereby informing more targeted implementation strategies for different player populations.

In this systematic review with meta-analysis, we aimed to examine the chronic effects of the FIFA 11 + on measures of physical fitness in male and female soccer players across different ages and performance levels, with specific fitness components identified on the basis of the outcomes reported in the included studies. Findings from this study will not only update the available knowledge base on the effectiveness of FIFA 11 + but also provide actionable insights for coaches, trainers and sports scientists seeking to enhance soccer performance through improvements in dynamic balance, agility, neuromuscular control and injury resilience.

2 Methods

This systematic review was carried out in accordance with the recommendations provided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses [32], and it was filed in advance in the PROSPERO register for systematic reviews (reference: CRD42025633810).

2.1 Literature Search Strategy

A computerized final literature search was conducted on 13 April, 2025 (Initial search conducted in 2025 and updated on 13 April, 2025) with two independent reviewers (I.N.K.V. and K.G.) systematically evaluating the literature searches. The search was performed across three major databases: MEDLINE (PubMed), Web of Science and Scopus [33]. To ensure a comprehensive review, keywords were selected on the basis of expert opinions and an extensive evaluation of the relevant literature [34]. Manual searches of the reference lists of earlier published systematic reviews were also conducted to locate other papers [35]. Boolean logic was employed to structure search queries, incorporating various keyword phrases, MeSH terms and combinations thereof. The keywords utilized for the search syntax included the following: 'FIFA 11 +,' 'FIFA 11 plus,' 'FIFA 11 + program*,' '11 +,' 'FIFA 11 plus program*,' 'football player*,' 'soccer player*,' 'football,' 'soccer,' 'speed,' 'sprint,' 'agility,' 'change of direction,' 'change-of-direction,' 'COD,' 'balance,' 'jump,' 'plyometric*,' 'motor skills,' 'muscle power' and 'athletic performance.' These terms were carefully selected using the operators AND and OR to capture the breadth of research relevant to the study's focus. Table 1 provides a detailed summary of the three distinct database search strategies employed in the review.

Table 3 Subgroup characteristics

Study	Duration	Frequency /week	Age	Sex	Tier level
Akbari et al. [63]	<9 weeks	≥ 3 sessions	< 18 years	Male	4
Arede et al. [10]	<9 weeks	< 3 sessions	< 18 years	Male	3
Asgari et al. [54]	≥ 9 weeks	≥ 3 sessions	< 18 years	Male	2
Ayala et al. [55]	<9 weeks	≥ 3 sessions	< 18 years	Male	2
Costa Silva et al. [59]	≥ 9 weeks	≥ 3 sessions	≥ 18 years	Male	3
Daneshjoo et al. [56]	<9 weeks	< 3 sessions	≥ 18 years	Male	2
Daneshjoo et al. [13]	<9 weeks	≥ 3 sessions	≥ 18 years	Male	2
Foqha et al. [53]	≥ 9 weeks	≥ 3 sessions	≥ 18 years	Male	3
Hwang et al. [64]	≥ 9 weeks	≥ 3 sessions	≥ 18 years	Male	2
Impellizzeri et al. [61]	≥ 9 weeks	≥ 3 sessions	≥ 18 years	Male	3
Nawed et al. [52]	≥ 9 weeks	≥ 3 sessions	≥ 18 years	Male	3
Pardos-Mainer et al. [11]	≥ 9 weeks	< 3 sessions	< 18 years	Female	2
Parsons et al. [12]	≥ 9 weeks	< 3 sessions	< 18 years	Female	2
Robles-Palazón et al. [58]	<9 weeks	≥ 3 sessions	< 18 years	Male	2
Sharifatpour et al. [9]	≥ 9 weeks	≥ 3 sessions	≥ 18 years	Male	3
Zarei et al. [57]	≥ 9 weeks	< 3 sessions	< 18 years	Male	2
Zhou et al. [62]	<9 weeks	≥ 3 sessions	< 18 years	Male	2

Tier level 2 = trained/developmental, Tier level 3 = highly trained/national level, Tier level 4 = elite/international level

Table 4 Study characteristics

Study	Country	<i>n</i>	Age (years)	Players (level/ sex)	Duration (week)	Intervention/ duration of intervention per session	Comparator	F/w	Specific physical fitness outcomes
Akbari et al. [63]	Iran	FG = 12 CG = 12	15–18	Elite soccer players	8	FIFA 11 + /25 min	Regular warm-up	3	Vertical jump height (VJ) ↑*
Arede et al. [10]	Portugal	FG = 15 CG = 15	10–14	Young soccer players	4	FIFA 11 + /20 min	Regular warm-up	3	Speed (20 m ↓*) CoD speed (T-test) ↓* Balance (YBT) ↑* Vertical jump height (VJ and CMJ) ↑*
Asgari et al. [54]	Iran	FG = 29 CG = 30 MG = 31	16–18	Youth football players	16	FIFA 11 + /25 min	Regular warm-up	3	CoD speed (IAT) ↓*
Ayala et al. [55]	Spain	FG = 10 CG = 11	16–18	Youth football players	4	FIFA 11 + /25 min	Regular warm-up	3	Speed (10 ↑ and 20 m ↓*) CoD speed (IAT) ↓* Balance (YBT) ↑* Vertical jump height (VDJ) ↑*
Costa Silva et al. [59]	Brazil	FG = 10 CG = 10	16–20	Junior soccer players	9	FIFA 11 + /20 min	Normal warm- up	3	Vertical jump height (CMJ and SJ) ↑*
Daneshjoo et al. [56]	Malaysia	FG = 12 CG = 12 HG = 12	18–22	Soccer players	8	FIFA 11 + /20 min	Regular warm-up	3	Balance (SEBT) ↑*
Daneshjoo et al. [13]	Malaysia	FG = 12 CG = 12 HG = 12	17–20	Young soccer players	8	FIFA 11 + /20 min	Regular warm-up	3	Speed (10 m and 20 m) ↓* CoD speed (IAT) ↓* Vertical jump height (VJ) ↑*
Foqha et al. [53]	Tunisia	FG = 13 CG = 12	24–29	Seven-a-side Soccer Athletes	10	FIFA 11 + /20 min	Regular warm-up	3	Speed (10 ↓* and 20 m ↑) CoD speed (MAT) ↓*
Hwang et al. [64]	South Korea	FG = 10 CG = 10	19–21	Collegiate soccer play- ers	12	FIFA 11 + /25 min	Normal warm- up	5	CoD speed (AAT-R and L) ↓* Speed (10 and 30-m ↓*)
Impellizzeri et al. [61]	Italy	FG = 42 CG = 39	19–28	Football play- ers	9	FIFA 11 + /25 min	Regular warm-up	3	Speed (20 m) ↓ CoD speed (T-Test) ↓ Balance (SEBT) ↑ Vertical jump height (VJ) ↑
Nawed et al. [52]	India	FG = 29 CG = 28	18–23	Soccer players	12	FIFA 11 + /25 min	Regular warm-up	5	Speed (20 m) ↓* CoD speed (T-test, and IAT) ↓ Vertical jump height (VJ) ↑*

Table 4 (continued)

Study	Country	<i>n</i>	Age (years)	Players (level/ sex)	Duration (week)	Intervention/ duration of intervention per session	Comparator	F/w	Specific physical fitness outcomes
Pardos-Mainer et al. [11]	Spain	FG = 19 CG = 17	12–14	Adolescent soccer play- ers	10	FIFA 11 +/15 min	Standard warm-up	2	CoD speed (V-cut Test) ↑* Balance (YBT) ↑* Vertical jump height (CMJ) ↑*
Parsons et al. [12]	Canada	FG = 25 CG = 18	9–11	Youth soccer players	20 (5-month)	FIFA 11 +/30 min	Regular warm-up (Coach determined)	1–2	Balance (YBT) ↓ CoD speed (T-Test) ↓ Jump High (VJ)↑
Robles- Palazón et al. [58]	Spain	FG = 10 CG = 11	15–18	Youth football players	4	FIFA 11 +/25 min	Regular warm-up	3	Balance (YBT) ↑ Speed (10- m↑ and 20-m↓) Jump High (VJ)=
Sharifatpour et al. [9]	Iran	FG = 20 CG = 20	22–28	Beach soccer player	12	FIFA 11 +/–	General warm-up	3	Jump High (Bosco index) ↑*
Zarei et al. [57]	Iran	FG = 34 CG = 32	14–16	Adolescent soccer play- ers	30	FIFA 11 +/25 min	Standard warm-up	2	CoD speed (IAT) ↓* Speed (9.1-m and 36.6 m) ↓* Jump High (VJ) ↑*
Zhou et al. [62]	China	FG = 10 CG = 10	10–15	Youth soccer players	8	FIFA 11 +/20 min	Regular warm-up	3	Speed (30 m) ↓* CoD speed (SR) ↓*

YBT Y balance test, *VJ* vertical jump, *MAT* modified agility test, *IAT* Illinois agility test, *SR* Shuttle run test, *SEBT* star excursion balance test, *VJ* vertical jump, *CMJ* countermovement jump, *AAT* arrow agility test, *SJ* squat jump, *CoD* change of direction, ↑ Value increased, ↑* Value significantly increased, ↓ Value decreased, ↓* Value significantly decreased, *FG* FIFA 11 + group, *CG* control group, *MG* modified FIFA 11 + group, *HG* HarmoKnee group, – not mentioned, *F/w* frequency per week, *n* number of participant in each group

2.2 Inclusion and Exclusion Criteria

To ensure a systematic and rigorous selection of studies, the Participants, Intervention, Comparators, Outcomes, and Study (PICOS) design framework was utilized [36]. This approach provided a structured method for evaluating study eligibility on the basis of clearly defined criteria. The inclusion and exclusion criteria adopted in this meta-analysis were designed to identify high-quality studies relevant to the objectives, while excluding research that did not meet the necessary standards. For the purpose of this review, studies that fulfilled the criteria are presented in Table 2.

2.3 Selection Process

Relevant data from the included studies were extracted into Microsoft Excel templates containing the titles of the articles and their corresponding DOIs. After removing duplicate records, the remaining documents were tabulated for further processing. Preliminary inclusion based on the PICOS framework (Population, Intervention, Comparison, Outcome and Study Design) was assessed collaboratively by all authors. This initial screening ensured that the studies met the predefined eligibility criteria before proceeding to detailed data extraction. Two authors (I.N.K.V. and K.G.) independently extracted key details




Fig. 2 Representation of Cochrane Risk of Bias Tool-1 (ROB1) summary: **a** ROB1 for individual studies, **b** Overall ROB1

from the included studies. These details comprised the first author's last name, year of publication, country of origin, study design, sample size, intervention specifics (duration, frequency and follow-up), outcome measures

and study findings. A third author (G.V.) verified the data for accuracy and completeness, providing an additional layer of validation [37]. Figure 1 shows the flowchart that depicts the study selection procedure for meta-analysis.




Fig. 3 Forest plot on the effects of FIFA 11+ versus active controls on measures of balance in soccer players




Fig. 4 Forest plot on the effects of FIFA 11+ versus active controls on measures of change-of-direction speed in soccer

2.4 Risk of Bias Assessment

The quality of the included randomized controlled trials (RCTs) was evaluated by two independent reviewers (I.N.K.V. and K.G.) using the Cochrane Risk of Bias Tool (ROB 1) [35]. To assess the Cochrane risk of bias, we focused on five domains: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting and other bias [38]. In addition, two authors (G.V. and M.E.) independently reviewed the risk of bias for each included study, with any disagreements

resolved through discussion and consensus involving a third and last author (G.V. and U.G.). Funnel plots were additionally created to identify potential publication bias (Electronic Supplementary Material [2]).

2.5 Data Extraction

Data pertaining to physical fitness, including measures of muscle power, linear sprint and CoD speed, agility and balance [39], were retrieved from each included research article. Dynamic balance was assessed using tests such as the Y balance test (YBT) or the star excursion balance test




Fig. 5 Forest plot on the effects of FIFA 11+ versus active controls on measures of linear sprint a 10-m and b 20-m in soccer players




Fig. 6 Forest plot on the effects of FIFA 11+ versus active controls on measures of muscle power (vertical jump height) in soccer

(SEBT). As proxies of muscle power, vertical jump height was assessed as indicated by the countermovement jump (CMJ), the drop jump (VDJ), the squat jump (SJ) test and the Bosco index. CoD speed and agility were evaluated through the T-Test, the Illinois Agility Test (IAT), the modified agility test (MAT), the arrow agility test (AAT) and the V-cut test. Linear sprint speed was assessed using the 10-m, 20-m and 30-m sprint tests. One author (I.N.K.V.) extracted the means, standard deviations (SD) and sample sizes (n) from

the publications that were included in the study, and a second author (K.G.) verified the accuracy of the retrieved data. The results were presented in the Electronic Supplementary Material [1]. In the event that there was a disagreement between the researchers, it was settled by conversation with a fourth author (M.E.). These characteristics included the participant's age, total duration of the intervention, training frequency, duration of the training session, participant's age, type of training protocol and the outcome measures that

Table 5 Subgroup analyses of potential moderators on balance, speed (10- and 20-m linear sprint and change of direction speed), muscle power (vertical jump height)

	Group (n)	# of studies	SMDs (95% CI)	I ²	p value	p _{diff}
<i>Balance outcomes</i>						
<i>Duration</i>						
≥ 9 weeks	6	3	0.17 (−0.07, 0.41)	0	0.15	0.02
< 9 weeks	9	4	0.62 (0.34, 0.90)	0	<0.001	
<i>Training frequency</i>						
≥ 3 sessions	7	3	0.47 (0.18, 0.75)	1	0.001	0.38
< 3 sessions	8	4	0.30 (0.04, 0.55)	14	0.02	
<i>Session duration</i>						
≥ 25 min	9	4	0.34 (0.07, 0.62)	21	0.01	0.61
< 25 min	6	3	0.45 (0.16, 0.74)	0	0.002	
<i>Age</i>						
≥ 18 years	2	2	0.56 (−0.34, 1.45)	71	0.22	0.09
< 18 years	13	5	0.36 (0.16, 0.57)	0	<0.001	
<i>Sex</i>						
Male	10	5	0.49 (0.26, 0.73)	0	<0.001	0.41
Female	5	2	0.17 (−0.11, 0.46)	0	0.23	
<i>Training and performance calibre</i>						
≥ Tier 3	3	2	0.26 (−0.07, 0.59)	0	0.13	0.41
< Tier 3	12	5	0.43 (0.18, 0.68)	21	<0.001	
<i>Change of direction speed outcomes</i>						
<i>Duration</i>						
≥ 9 weeks	10	8	−0.61 (−1.13, −0.08)	85	0.02	0.77
< 9 weeks	4	4	−0.80 (−1.93, 0.34)	84	0.17	
<i>Training frequency</i>						
≥ 3 sessions	10	8	−1.05 (−1.68, −0.42)	86	0.001	0.001
< 3 sessions	4	4	0.12 (−0.18, 0.42)	0	0.44	
<i>Session duration</i>						
≥ 25 min	9	7	−0.32 (−0.70, 0.06)	71	0.10	0.07
< 25 min	5	5	−1.72 (−3.21, −0.24)	92	0.02	
<i>Age</i>						
≥ 18 years	7	5	−1.45 (−2.37, −0.53)	91	0.002	0.004
< 18 years	7	7	−0.06 (−0.30, 0.18)	4	0.63	
<i>Sex</i>						
Males	12	10	−0.79 (−1.33, −0.25)	86	0.004	0.04
Females	2	2	−0.04 (−0.49, 0.40)	0	0.84	
<i>Training and performance calibre</i>						
≥ Tier 3	5	4	−1.00 (−2.04, 0.03)	92	0.06	0.41
< Tier 3	9	8	−0.53 (−1.00, −0.06)	73	0.03	
<i>10-m linear sprint speed outcomes</i>						
<i>Duration</i>						
≥ 9 weeks	3	3	0.40 (0.03, 0.78)	0	0.04	0.03
< 9 weeks	3	3	−1.17 (−2.53, 0.20)	83	0.10	
<i>Session duration</i>						
≥ 25 min	4	4	−0.24 (−1.23, 0.75)	84	0.64	0.64
< 25 min	2	2	−0.74 (−2.59, 1.11)	89	0.42	
<i>Age</i>						
≥ 18 years	3	3	−0.48 (−1.61, 0.65)	80	0.40	0.88
< 18 years	3	3	−0.35 (−1.74, 1.04)	89	0.62	
<i>20-m linear sprint speed</i>						
<i>Duration</i>						
≥ 9 weeks	3	3	−0.39 (−2.68, 1.90)	97	0.74	0.79

Table 5 (continued)

	Group (n)	# of studies	SMDs (95% CI)	I^2	p value	p_{diff}
<9 weeks	4	4	−0.72 (−1.51, 0.07)	70	0.07	
<i>Session duration</i>						
≥25 min	4	4	−0.93 (−2.20, 0.34)	92	0.15	0.53
<25 min	3	3	−0.16 (−2.24, 1.93)	94	0.88	
<i>Age</i>						
≥18 years	4	4	−0.82 (−2.68, 1.04)	96	0.39	0.60
<18 years	3	3	−0.31 (−0.77, 0.16)	0	0.19	
<i>Training and performance calibre</i>						
≥Tier 3	4	4	−0.36 (−1.96, 1.23)	95	0.65	0.58
<Tier 3	3	3	−0.91 (−2.02, 0.20)	77	0.11	
<i>Vertical jump height outcome</i>						
<i>Duration</i>						
≥9 weeks	9	7	0.26 (0.05, 0.46)	78	0.01	0.007
<9 weeks	6	5	0.79 (0.46, 1.13)	0	<0.001	
<i>Training frequency</i>						
≥3 sessions	9	8	0.73 (0.49, 0.96)	56	<0.001	<0.001
<3 sessions	6	4	0.01 (−0.25, 0.27)	63	0.94	
<i>Session duration</i>						
≥25 min	7	7	0.29 (0.06, 0.52)	68	0.01	0.45
<25 min	7	4	0.43 (0.14, 0.73)	74	0.004	
<i>Age</i>						
≥18 years	6	5	0.64 (0.37, 0.90)	68	<0.001	0.02
<18 years	9	7	0.22 (−0.02, 0.45)	70	0.07	
<i>Sex</i>						
Male	12	10	0.54 (0.34, 0.74)	71	<0.001	0.003
Female	3	2	−0.09 (−0.46, 0.28)	0	0.63	
<i>Training and performance calibre</i>						
≥Tier 3	8	6	0.75 (0.52, 0.99)	58	<0.001	<0.001
<Tier 3	7	6	−0.01 (−0.26, 0.25)	54	0.97	

SMD standardized mean difference, I^2 study heterogeneity, p_{diff} significant differences among two subgroups

Bold values indicate statistically significant subgroup differences ($p_{diff} < 0.05$)

were utilized in the study. These characteristics are presented in Table 3.

2.6 Statistical Analyses

The Review Manager Software (RevMan 5.4) was used to meta-analyse our findings [40]. The threshold for statistical significance was established at $p < 0.05$. The inverse-variance random-effects model was employed, which assigns weights to each trial inversely proportional to the variance of their effect estimates, thereby accounting for both within-study and between-study variability [41]. This approach facilitates a more generalizable estimate of the overall effect by incorporating potential heterogeneity across studies. The standardized mean differences (SMDs) alongside confidence intervals (95% CI) were used to describe the

magnitude of the observed effects. The SMD was interpreted using the following thresholds: <0.2 trivial; 0.2–0.6, small; 0.6–1.2, moderate; 1.2–2.0, large; 2.0–4.0, very large; >4.0, extremely large [42–44]. To evaluate study heterogeneity among the group and subgroup analyses, heterogeneity (I^2) statistics were computed, with values categorized as follows: <25% indicating low heterogeneity, 25–75% representing moderate heterogeneity, and >75% reflecting high heterogeneity [45, 46]. Only outcome variables assessed in a minimum of five independent studies were included in the meta-analysis to ensure adequate statistical power, enhance the reliability of the findings and minimize the risk of drawing conclusions from limited evidence [47, 48].

Subgroup analyses were conducted to explore factors potentially moderating the effectiveness of FIFA 11+. Programming variables were categorized using the median split method, as outlined in prior research

[49], to facilitate meaningful comparison across studies, as it provides a straightforward and consistent way to classify continuous variables into balanced groups. The median was calculated if at least three studies provided data for a given moderator [50]. The following training-related factors were included: training duration (≥ 9 weeks versus < 9 weeks), weekly training frequency (≥ 3 sessions versus < 3 sessions), training and performance calibre ($\geq$ Tier 3 versus $<$ Tier 3), duration of a single training session (≥ 25 min versus < 25 min). The Participant Classification Framework (based on the training and performance calibre) was applied to define tier levels consistently, whereby Tier 1 is recreationally active, Tier 2 trained/developmental, Tier 3 highly trained/national level and Tier 4 elite/international level [51]. Age was classified on the basis of developmental stages, distinguishing between adults (≥ 18 years) and youth (< 18 years), while sex was categorized as male and female. These analyses were undertaken only when at least two studies were available in each subgroup to ensure meaningful comparisons. This approach provided insights into the consistency of the findings and the potential impact of specific variables on the intervention outcomes, ensuring a robust understanding of the results.

3 Results

The initial search identified 4829 articles from three databases and 18 from other sources (e.g., ResearchGate, Citations, Websites). After removing 1702 duplicates, 3127 articles remained for eligibility assessment. Title and abstract screening excluded 35 review articles and 2997 unrelated studies. This left 95 database-sourced articles and 18 from other sources for further evaluation. Full-text screening led to the exclusion of 96 articles on the basis of predefined PICOS criteria. Ultimately, 17 studies met the eligibility criteria and were included in the meta-analysis, as illustrated in Fig. 1. Notably, compared with the previous review [24], an additional nine studies that were either not included earlier or have been recently published were incorporated in the present analysis, thereby extending and updating the available evidence base.

3.1 Study Characteristics

The study characteristics are presented in Table 4. The included studies reported training durations ranging from 4 to 30 weeks, with training frequencies varying between 1 and 5 weekly sessions. Player's ages ranged from 9 to 29 years. A total of 611 soccer players were analysed in this meta-analysis, comprising 532 males (15 studies)

and 79 females (two studies). Soccer players' training and performance calibre ranged from Tier 2 to Tier 4, with 334 soccer players being rated as $<$ Tier 3 (seven studies), $277 \geq$ Tier 3 (ten studies). A total of 16 articles reported intervention durations ranging from 15 to 30 min per session. At the same time, one study did not specify the duration details [9]. Eight studies were classified as ≥ 18 years and nine as < 18 years. By training and performance calibre, seven studies were $\geq$ Tier 3 and ten $<$ Tier 3. These subgroup analyses highlight the impact of demographic and training programming variables on FIFA 11 + effectiveness, as shown in Table 3.

3.2 Methodological Quality of the Included Studies

The studies demonstrate varying levels of bias across different domains (Fig. 2). Random sequence generation was assessed as having a low risk of bias in all the included studies [9–13, 52–64]. Allocation concealment was frequently rated as unclear or high risk, reflecting insufficient reporting or inadequate methods. The blinding of participants and personnel presented the highest proportion of high-risk ratings, highlighting significant shortcomings in implementing blinding protocols. In contrast, blinding of outcome assessment generally showed a high risk of bias [9, 10, 12, 13, 52, 54, 55, 57, 59, 62–64], except for a few studies [56, 58, 61]. A total of 17 studies were included in the meta-analysis, with the overall risk of bias assessed using the ROB 1 tool, as illustrated in Fig. 2.

3.3 Effects of FIFA 11 + on Measures of Balance

A statistically significant difference was observed in favour of the FIFA 11 + for dynamic balance (seven studies: small SMDs = 0.37, CI 0.18–0.56, $p < 0.001$, $I^2 = 7$) (Fig. 3).

3.4 Effects of FIFA 11 + on Linear Sprint and Change-of-Direction Speed

FIFA 11 + induced significantly larger CoD speed improvements than active controls were more effective (12 studies: moderate SMDs = -0.65 , CI -1.11 , -0.20 , $p = 0.005$, $I^2 = 84$) (Fig. 4). No statistically significant difference was observed between the FIFA 11 + group and the control group for 10-m linear sprint speed (six studies: small SMDs = -0.40 , CI -1.22 to 0.42 , $p = 0.34$, $I^2 = 84$) (Fig. 5a) and 20-m linear sprint speed (seven studies:

moderate SMDs = 0.61, CI 1.61 to 0.40, $p = 0.24$, $I^2 = 92$) (Fig. 5b).

3.5 Effects of FIFA 11+ on Proxies of Muscle Power

Vertical jump height was assessed as a proxy of muscle power. A statistically significant difference was observed between the FIFA 11+ group and the active control group for vertical jump height (12 studies: small SMDs = 0.56, CI 0.22–0.89, $p < 0.001$, $I^2 = 71$) (Fig. 6).

The overall analysis on outcome measures demonstrated low-to-high study heterogeneity, with I^2 values ranging between 7 and 92%. Specifically, dynamic balance showed low heterogeneity ($I^2 = 7\%$), vertical jump height reflected moderate heterogeneity ($I^2 = 71\%$), while change-of-direction speed ($I^2 = 84\%$) and linear sprint outcomes indicated high heterogeneity (10-m sprint: $I^2 = 84\%$; 20-m sprint: $I^2 = 92\%$). These findings suggest variability in the treatment effects across the included studies, as illustrated in Figs. 3, 4, 5 and 6.

3.6 Subgroup Analyses According to Programming and Demographic Parameters

For training duration (≥ 9 versus < 9 weeks), larger FIFA 11+ effects were found for < 9 weeks. With regards to weekly training frequency (≥ 3 versus < 3 session), larger FIFA 11+ effects were found for ≥ 3 sessions. With regards to chronological age (≥ 18 versus < 18 years), larger FIFA 11+ effects were found for ≥ 18 years (i.e., adult soccer players). In terms of sex (males versus females), larger FIFA 11+ effects were found for male soccer players. For training and performance calibre ($\geq$ Tier 3 versus $<$ Tier 3), a larger effect was found for $\geq$ Tier 3 soccer players (Table 5).

4 Discussion

In this meta-analysis, we aimed to synthesize evidence from 17 studies on the effects of the FIFA 11+ program on balance, and CoD speed, muscle power (jump performance) in youth and adult soccer players. Notably, authors from all included studies reported at least one positive or beneficial effect associated with the FIFA 11+ program. The results of the meta-analysis revealed that the FIFA 11+ program produced small-to-moderate improvements in dynamic balance, linear sprint and CoD speed and jump performance when compared with conventional warm-up programs (general, standard or normal warm-ups). More specifically, for measures of dynamic balance, statistically significant but small effect sizes ($p < 0.001$) were found in favour of FIFA 11+. CoD speed improved significantly ($p < 0.01$) with moderate

effect sizes, while small effects were found for vertical jump height improvements ($p < 0.001$), all in favour of FIFA 11+. In contrast, linear sprint speed showed no statistically significant improvements following the application of FIFA 11+.

The observed improvements in dynamic balance following the FIFA 11+ intervention may be attributed to the specific exercises included in part two of the program. This section incorporates progressive plyometric, balance and core muscle exercises, all of which are designed to enhance neuromuscular control and postural stability [65]. These exercises likely contribute to improved proprioception, postural stability and coordination, which are essential components of dynamic balance. Thus, the positive outcomes observed in this meta-analysis may be closely related to the neuromuscular adaptations elicited by these targeted training elements during the intervention [52]. Few systematic reviews have reported a positive effect on dynamic balance following various warming-up programs [7, 66, 67]. Within the sport of soccer, the ability to change directions is among the most crucial components [68]. Change of direction (CoD) ability is a multifaceted skill that relies on speed, coordination, dynamic balance, strength and reactive agility [69]. The improvements in CoD speed observed in this study may be attributed to the specific exercises in parts two and three of the FIFA 11+ program; part one includes progressive plyometric, balance and strength-based drills designed to enhance neuromuscular coordination and control. Part 2 includes different running exercises (cross-pitch running, bounding and running, plant and cut running) [65]. These exercises target essential movement qualities, such as rapid deceleration, re-acceleration and stabilization, which are critical for CoD performance [17, 70]. There has been scant evidence from systematic reviews that different warming-up programs improve CoD [7]. Although these findings support the efficacy of FIFA 11+ for improving CoD speed and balance, caution is warranted in generalizing these results to all soccer-playing populations. Most included studies featured male athletes from specific age ranges and competitive/Tier levels [9, 52, 54, 56, 57, 63, 64], and there is limited evidence on female players [11, 12], youth athletes [10, 58, 60, 62] or professional players [13, 53, 63]. Consequently, the observed performance benefits may not directly translate across all demographic and competitive groups, underscoring the need for broader, more representative research. The observed improvements in vertical jump performance following the FIFA 11+ intervention may be attributed to the progressive plyometric and strength exercises included in part two of the program. These exercises, such as planks, side planks, Nordic hamstring drills, single-leg balance activities, squats and various jumping drills (vertical, lateral and box jumps), are progressively structured to enhance core stability, lower-limb strength and explosive power [17]. By targeting neuromuscular coordination, reactive

ability and biomechanical efficiency, these exercises play a crucial role in optimizing jump performance. Thus, the positive outcomes in vertical jump height observed in this study are likely a result of the neuromuscular adaptations induced by these training elements during the intervention. A small number of systematic evaluations have documented improvements in jump performance after different warm-up programs [71, 72]. However, it is important to note that such outcomes may also depend on intervention fidelity. Variability in adherence, supervision and delivery of the FIFA 11+ across studies likely influenced the magnitude of improvements observed. While some investigations ensured structured and consistent implementation, others applied it less rigorously, which may have contributed to heterogeneity in effects. These differences in fidelity and compliance should therefore be considered when interpreting pooled results [73, 74].

Subgroup analyses revealed several significant findings regarding the effects of the FIFA 11+ intervention on balance, change of direction, speed performance and jump performance among soccer players. In terms of training duration, shorter programs (<9 weeks) significantly enhanced CoD speed and jump performance versus longer programs (≥ 9 weeks), possibly owing to their initial responsiveness to new training stimuli, whereas longer durations may lead to diminishing returns as players adapt to the regimen [75]. Training frequency played a crucial role in determining outcomes, with players training ≥ 3 sessions per week exhibiting superior improvements in CoD and vertical jump height, likely owing to consistent exposure to the intervention [76]. Interestingly, players training <3 sessions per week achieved better results in 10 m speed, potentially because reduced frequency allowed for better recovery and adaptation specific to this metric [77]. Soccer players aged ≥ 18 years demonstrated greater improvements in CoD speed and jump performance versus those aged <18 years, likely owing to the more advanced neuromuscular and physiological maturity of the older group, which enabled better adaptation to the training program [21, 78]. This suggests that maturity of the neuromuscular system, characterized by more efficient motor unit recruitment, enhanced coordination and improved muscle-tendon stiffness in older players [78], may provide a biological advantage in responding to structured interventions such as FIFA 11+.

The subgroup analyses demonstrated that male versus female soccer players exhibited significantly greater improvements in CoD speed and vertical jump height, suggesting a higher neuromuscular response to the FIFA 11+ program. This may be attributed to sex-related physiological differences, such as greater muscle mass and power output in males [79]. However, sex-specific sports participation patterns may also play a role,

as prior evidence indicates that female athletes often face contextual barriers (e.g., lower training adherence, fewer structured exposure hours) that could attenuate potential benefits of the training program, despite its injury-preventive and performance-enhancing capacity [80, 81]. In addition, higher training and performance calibre were associated with greater vertical jump height, with $\geq$ Tier 3 players showing significantly larger gains compared with < Tier 3. Notably, only one study fell under Tier 4 [63], while seven studies were classified as Tier 3. Higher-calibre players likely undergo more intensive and sport-specific training, leading to superior neuromuscular efficiency, enhanced motor unit recruitment and increased rate of force development, all of which contribute to improved maximal power production [82]. These findings may also reflect higher baseline fitness levels in elite players, which facilitate greater responsiveness to structured warm-up and plyometric components embedded within FIFA 11+, as they can more effectively translate training stimuli into performance gains [61, 63]. Vertical jump height, reflecting athletes' higher baseline fitness, advanced neuromuscular efficiency and greater ability to capitalize on structured plyometric training, included interventions such as the FIFA 11+ program [24, 83]. These results underscore the importance of considering factors such as training duration, frequency, age and competitive level when designing and implementing FIFA 11+ programs to improve players' physical fitness. In addition, our findings highlight that the effects of the FIFA 11+ program may not only vary across moderators such as age, training and performance calibre but also depend on whether adaptations are acute (short-term neuromuscular responses) or chronic (long-term physiological changes). This distinction may help explain the heterogeneity observed across studies and reinforces the program's versatility for both acute performance potentiation and sustained athletic development. Moreover, our findings suggest that the program's effects may differ across player groups. Tailored adaptations for female athletes, considering sex-specific neuromuscular and physiological characteristics, and for youth athletes, aligned with developmental stages, could further enhance its effectiveness and ensure safe application.

4.1 Limitations and Future Recommendations

Several limitations and recommendations for future research agendas must be acknowledged. First of all, there is a significant lack of studies on the use of the FIFA 11+ program among female soccer players. Accordingly, the current findings, mostly from male soccer players, cannot be generalized to female populations, emphasizing the necessity for research into the program's impact on female athletes to close this gap in the literature. In addition, underrepresented populations, including youth and amateur players, should be included to improve the generalizability of findings. Furthermore, future research should strive

to report more information on programming variables such as training frequency, intensity and volume, but also demographic variables including sex, chronological and biological age. By providing detailed information on these relevant issues, FIFA 11 + programs can be better targeted to the needs of the respective population under investigation. Moreover, high-quality, low-bias RCTs with standardized implementation of FIFA 11 + are needed to strengthen causal inferences. Reporting standardized outcome measures and effect sizes alongside SMDs would help clarify the clinical significance of performance improvements. Finally, many included studies lacked robust methodological rigour, particularly in terms of allocation, concealment and blinding processes. These shortcomings may cause bias that could affect the robustness of findings. Future studies should prioritize rigorous experimental designs with proper allocation concealment and blinding to enhance the credibility and generalizability of the results. In addition, heavy reliance on SMDs and high heterogeneity in certain outcomes, such as sprint performance, may limit confidence in some findings. Future research with larger sample sizes should conduct sensitivity analyses and subgroup assessments to strengthen the stability of results. In addition, although high heterogeneity was observed in the sprint outcomes (10-m and 20-m), no sensitivity analysis was conducted owing to the absence of statistically significant effects and the limited number of studies available for these measures, which could have compromised the stability of further subgroup analyses. This should be addressed in future research with larger datasets to enable more robust heterogeneity assessments.

4.2 Practical Implications

This meta-analysis highlights the practical value of incorporating the FIFA 11 + program into routine soccer warm-ups to enhance key physical fitness attributes in both youth and adult players. Coaches, strength and conditioning professionals and physical trainers are encouraged to replace or supplement conventional warm-ups with the FIFA 11 + to improve dynamic balance, change-of-direction speed and vertical jump performance. While optimal benefits were observed when the program was performed at least three times per week and over shorter intervention periods (<9 weeks), achieving such adherence in real-world settings may be challenging owing to factors such as match schedules, travel demands and overall training load. Therefore, practitioners should consider flexible implementation strategies, such as integrating selected FIFA 11 + exercises into technical or tactical drills, to maintain regularity. Given the substantial heterogeneity across studies, the program's effectiveness may vary depending on age, competitive level and baseline physical characteristics, with greater effects observed in older, male and higher-tier players. However, the underrepresentation of female participants limits the generalizability of these findings to women's soccer, highlighting the need for further research in more diverse and representative populations.

When implemented with a context-specific approach, the FIFA 11 + can serve as an effective tool for both performance enhancement and injury prevention, supporting long-term athlete development and competitive success.

5 Conclusions

This meta-analysis concluded from 17 studies, significant chronic effects of the FIFA 11 + program versus conventional soccer-specific warm-up programs on key physical fitness components, including dynamic balance, and CoD speed, as well as vertical jump performance. The observed benefits, while statistically significant, were generally of small-to-moderate magnitude and may have been influenced by variability across studies. The findings highlight the potential of FIFA 11 + to induce neuromuscular adaptations in youth and adult soccer players, resulting in improved linear sprint and CoD speed as well as muscle strength and power, important fitness qualities related to success in soccer. This study contributes to the broader field of sports science by providing robust evidence of the program's capacity to optimize physical performance, offering valuable insights for athletes, coaches and sports organizations globally. However, the magnitude of improvement should be interpreted in light of the heterogeneity and differences in program implementation in the study. By establishing a foundation of scientifically validated benefits, the research encourages the integration of structured, evidence-based training protocols into routine practice. The demonstrated effectiveness of the FIFA 11 + program supports its adoption across diverse sports disciplines, fostering enhanced performance levels and setting new benchmarks for athletic excellence. This global perspective underscores the program's transformative potential to redefine training strategies and contribute to the sustained growth and development of athletic performance worldwide.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s40279-025-02346-8>.

Funding Open Access funding enabled and organized by Projekt DEAL.

Declarations

Funding We acknowledge support by the Open Access Publication Fund of the University of Freiburg, Germany.

Ethics approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

Code availability Not applicable.

Conflict of interest The authors declare they that have no conflicts of interest with the content of this article. Urs Granacher is a member of the editorial board of Sports Medicine but was not involved in the selection of reviewers for this manuscript or in any of the subsequent editorial decisions.

Availability of data and material The data are provided in Table 4 and Electronic Supplementary Material [1], with further datasets available from the corresponding author upon reasonable request.

Authors' contribution Conceptualization: INKV, KG. Literature search: INKV, KG, GV. Abstract screening, full-text eligibility assessment, data extraction and critical appraisal: INKV, KG, GV, ME, HŞU, UG. Data synthesis: INKV, KG, CCTC, KP, BK, HZ, UG. Original draft of manuscript: INKV, KG, UG. Manuscript review and edit: INKV, CCTC, KP, BK, HŞU, HZ, UG. All authors read and approved the final manuscript.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Fradkin AJ, Zazryn TR, Smoliga JM. Effects of warming-up on physical performance: a systematic review with meta-analysis. *J Strength Cond Res*. 2010;24(1):140–8. <https://doi.org/10.1519/JSC.0b013e3181c643a0>.
2. Thorborg K, Krommes KK, Esteve E, Clausen MB, Bartels EM, Rathleff MS. Effect of specific exercise-based football injury prevention programmes on the overall injury rate in football: a systematic review and meta-analysis of the FIFA 11 and 11+ programmes. *Br J Sports Med*. 2017;51(7):562–71. <https://doi.org/10.1136/bjsports-2016-097066>.
3. Patti A, Giustino V, Cataldi S, Stoppa V, Ferrando F, Marvulli R, et al. Effects of 5-week of FIFA 11+ warm-up program on explosive strength, speed, and perception of physical exertion in elite female futsal athletes. *Sports*. 2022;10(7):100. <https://doi.org/10.3390/sports10070100>.
4. Vlachas T, Paraskevopoulos E. The effect of the FIFA 11+ on injury prevention and performance in football: a systematic review with meta-analysis. *BioMed*. 2022;2(3):328–40. <https://doi.org/10.3390/biomed2030026>.
5. Ayala F, Calderón-López A, Delgado-Gosálbez JC, Parra-Sánchez S, Pomares-Noguera C, Hernández-Sánchez S, et al. Acute effects of three neuromuscular warm-up strategies on several physical performance measures in football players. *PLoS One*. 2017;12(1):1–17. <https://doi.org/10.1371/journal.pone.0169660>.
6. Asgari M, Schmidt M, Terschluse B, Sueck M, Jaitner T. Acute effects of the FIFA11+ and Football+ warm-ups on motor performance. A crossover randomized controlled trial. *PLoS One*. 2023;18:1–13. <https://doi.org/10.1371/journal.pone.0284702>. (4 April).
7. Gomes Neto M, Conceição CS, De Lima Brasileiro AJA, De Sousa CS, Carvalho VO, De Jesus FLA. Effects of the FIFA 11 training program on injury prevention and performance in football players: a systematic review and meta-analysis. *Clin Rehabil*. 2017;31(5):651–9. <https://doi.org/10.1177/0269215516675906>.
8. Bizzini M, Junge A, Dvorak J. Implementation of the FIFA 11+ football warm up program: how to approach and convince the Football associations to invest in prevention. *Br J Sports Med*. 2013;47(12):803–6. <https://doi.org/10.1136/bjsports-2012-092124>.
9. Sharifatpour R, Abbasi H, Alizadeh MH, Abedinzadeh S, Machado S. Effect of 12 weeks of modified FIFA 11+ training on functional factors of male beach soccer players: a randomized controlled trial study. *Cuad Psicol Deporte*. 2024;24(3):265–80. <https://doi.org/10.6018/cpd.584601>.
10. Arede J, Shang X, Moran J, Calleja-González J, Travassos B, Madruga-Parera M, et al. Does the FIFA 11 + warm-up program positively shape both the physical and small-sided game performance of youth football players? *Intell Sport Heal*. 2025;1(1):22–9. <https://doi.org/10.1016/j.ish.2024.12.004>.
11. Pardos-Mainer E, Casajús JA, Gonzalo-Skok O. Adolescent female soccer players' soccer-specific warm-up effects on performance and inter-limb asymmetries. *Biol Sport*. 2019;36(3):199–207. <https://doi.org/10.5114/biolsport.2019.85453>.
12. Parsons JL, Carswell J, Nwoba IM, Stenberg H. Athlete perceptions and physical performance effects of the FIFA 11+ program in 9-11 year-old female soccer players: a cluster randomized trial. *Int J Sports Phys Ther*. 2019;14(5):740–52. <https://doi.org/10.26603/ijsp20190740>.
13. Daneshjoo A, Mokhtar AH, Rahnama N, Yusof A. Effects of the 11+ and Harmoknee warm-up programs on physical performance measures in professional soccer players. *J Sports Sci Med*. 2013;12(3):489–96.
14. Ferreira-Júnior JB, da Encarnação IGA, Rezende VHS, Oliveira JCG, Souza LA, Assunção JC, et al. Effects of different intensities of FIFA 11+ on physical performance of U-15 soccer players. *Apunts Sports Med*. 2023. <https://doi.org/10.1016/j.apunsm.2023.100419>.
15. Zarei M, Eshghi S, Hosseinzadeh M. The effect of a shoulder injury prevention programme on proprioception and dynamic stability of young volleyball players; a randomized controlled trial. *BMC Sports Sci Med Rehabil*. 2021;13(1):71. <https://doi.org/10.1186/s13102-021-00300-5>.
16. Sahin N, Gürses VV, Baydil B, Akgül MS, Feka K, Iovane A, et al. The effect of comprehensive warm up (fifa 11+ program) on motor abilities in young basketball players: a pilot study. *Acta Med Mediterr*. 2018;34(3):703–8. https://doi.org/10.19193/0393-6384_2018_3_108.
17. Cerrillo-Sanchis J, Muñoz-Criado I, Pérez-Puchades V, Palmero-Martín I, Galcerán-Ruiz J, Portes-Sanchez R, et al. Applying a specific warm-up on basketball performance: the basket-up approach. *Appl Sci*. 2024;14(14):5969. <https://doi.org/10.3390/app14145969>.
18. Singh VR, Stokes K, McKay CD. Feasibility and acceptability of a new shoulder-specific warm-up programme to prevent injuries in community youth rugby union as compared to the FIFA 11+. *Int J Sports Sci Coach*. 2024;19(3):1093–104. <https://doi.org/10.1177/17479541231192170>.
19. Attwood MJ, Roberts SP, Trewartha G, England ME, Stokes KA. Efficacy of a movement control injury prevention programme in adult men's community rugby union: a cluster

- randomised controlled trial. *Br J Sports Med.* 2018;52(6):368–74. <https://doi.org/10.1136/bjsports-2017-098005>.
20. Lopes M, Oliveira J, Ribeiro F. Injury prevention in futsal players: is the FIFA 11+ a simple answer to a complex problem? *Phys Ther Rev.* 2020;25(2):96–105. <https://doi.org/10.1080/10833196.2020.1731177>.
21. Markovic G, Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sports Med.* 2010;40(10):859–95. <https://doi.org/10.2165/11318370-000000000-00000>.
22. Isla E, Romero-Moraleda B, Moya J, Esparza-Ros F, Mallo J. Effects of a neuromuscular warm-up program in youth female soccer players. *J Hum Kinet.* 2021;79(1):29–40. <https://doi.org/10.2478/hukin-2021-0080>.
23. Granacher U, Behm DG, Faigenbaum AD. Neuromuscular training effects on performance. In: Armstrong N, van Mechelen W, editors. *Oxford Textbook of Children's Sport and Exercise Medicine* 4e. Oxford: Oxford University Press; 2023. p. 505–20. <https://doi.org/10.1093/med/9780192843968.003.0036>.
24. Liu R, Liu J, Ma X, Li Q, An R. Effect of FIFA 11+ intervention on change of direction performance in soccer and futsal players: a systematic review and meta-analysis. *Int J Sport Sci Coach.* 2021;16(3):862–72. <https://doi.org/10.1177/1747954121991667>.
25. Baroni BM, Couto W, Leal Junior ECP. Estudo descritivo-comparativo de parâmetros de desempenho aeróbio de atletas profissionais de futebol e futsal. *Rev Bras Cineantropom Desempenho Hum.* 2011;13(3):170–6. <https://doi.org/10.5007/1980-0037.2011v13n3p170>.
26. Burdukiewicz A, Pietraszewska J, Stachon A, Chromik K, Golin-ski D. The anthropometric characteristics of futsal players compared with professional soccer players. *Hum Mov.* 2014;15(2):93–9. <https://doi.org/10.2478/humo-2014-0008>.
27. Jiménez-Reyes P, García-Ramos A, Cuadrado-Peñañiel V, Párraga-Montilla JA, Morcillo-Losa JA, Samozino P, et al. Differences in sprint mechanical force–velocity profile between trained soccer and futsal players. *Int J Sports Physiol Perform.* 2019;14(4):478–85. <https://doi.org/10.1123/ijspp.2018-0402>.
28. Steffen K, Bakka HM, Myklebust G, Bahr R. Performance aspects of an injury prevention program: a ten-week intervention in adolescent female football players. *Scand J Med Sci Sports.* 2008;18(5):596–604. <https://doi.org/10.1111/j.1600-0838.2007.00708.x>.
29. Silva JR, Nassis GP, Rebelo A. Strength training in soccer with a specific focus on highly trained players. *Sports Med Open.* 2015;1(1):17. <https://doi.org/10.1186/s40798-015-0006-z>.
30. Lesinski M, Prieske O, Granacher U. Effects and dose-response relationships of resistance training on physical performance in youth athletes: a systematic review and meta-analysis. *Br J Sports Med.* 2016;50(13):781–95. <https://doi.org/10.1136/bjsports-2015-095497>.
31. Asgari M, Nazari B, Bizzini M, Jaitner T. Effects of the FIFA 11+ program on performance, biomechanical measures, and physiological responses: a systematic review. *J Sport Health Sci.* 2023;12(2):226–35. <https://doi.org/10.1016/j.jshs.2022.05.001>.
32. Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ.* 2015;349:g7647. <https://doi.org/10.1136/bmj.g7647>.
33. Ramirez-Campillo R, Perez-Castilla A, Thapa RK, Afonso J, Clemente FM, Colado JC, et al. Effects of plyometric jump training on measures of physical fitness and sport-specific performance of water sports athletes: a systematic review with meta-analysis. *Sports Med Open.* 2022;8(1):108. <https://doi.org/10.1186/s40798-022-00502-2>.
34. Gavagni VZ, Vahed N. Key word validity ratio (kwvr); a tool for validity of keywords for building an accurate literature search strategy. *BMJ Open.* 2017;7:184. <https://doi.org/10.1136/bmjopen-2016-015415.184>.
35. Boopalan D, Vijayakumar V, Ravi P, Shanmugam P, Kuppusamy M, Masilamani E, et al. Effect of yoga as a complementary therapy in prostate cancer survivors: a systematic review. *Support Care Cancer.* 2024;32(8):1–9. <https://doi.org/10.1007/s00520-024-08754-2>.
36. Higgins JPT, Green S, Ben Van Den A. *Cochrane handbook for systematic reviews of interventions.* *Int Coach Psychol Rev.* 2020;15(2):123–5. <https://doi.org/10.53841/bpsicpr.2020.15.2.123>.
37. Whiting P, Rutjes A, Dinnes J, Reitsma J, Bossuyt P, Kleijnen J. Development and validation of methods for assessing the quality of diagnostic accuracy studies. *Health Technol Assess.* 2004;8(25):1–234. <https://doi.org/10.3310/hta8250>.
38. Reveiz L, Chapman E, Asial S, Munoz S, Bonfill X, Alonso-Coello P. Risk of bias of randomized trials over time. *J Clin Epidemiol.* 2015;68(9):1036–45. <https://doi.org/10.1016/j.jclinepi.2014.06.001>.
39. Falces-Prieto M, González-Fernández FT, García-Delgado G, Silva R, Nobari H, Clemente FM. Relationship between sprint, jump, dynamic balance with the change of direction on young soccer players' performance. *Sci Rep.* 2022;12(1):1–9. <https://doi.org/10.1038/s41598-022-16558-9>.
40. Sun Y, Lu B, Su W, Song X, Shang X, Zheng J, et al. Comprehensive assessment of the effects of concurrent strength and endurance training on lipid profile, glycemic control, and insulin resistance in type 2 diabetes: a meta-analysis. *Medicine.* 2024;103(12):E37494. <https://doi.org/10.1097/MD.00000000000037494>.
41. Traise A, Dieberg G, Pearson MJ, Smart NA. The effect of exercise training in people with pre-dialysis chronic kidney disease: a systematic review with meta-analysis. *J Nephrol.* 2024;37(8):2063–98. <https://doi.org/10.1007/s40620-024-02081-9>.
42. Suso-Martí L, Núñez-Cortés R, Sánchez-Sabater A, Garrigós-Pedron M, Ferrer-Sargues FJ, López-Bueno R, et al. Effects of exercise-based interventions on inflammatory markers in patients with fibromyalgia: a systematic review and meta-analysis. *Semin Arthritis Rheum.* 2024;65:152377. <https://doi.org/10.1016/j.semarthrit.2024.152377>.
43. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3–12. <https://doi.org/10.1249/MSS.0b013e31818cb278>.
44. Garcia-Carrillo E, Ramirez-Campillo R, Izquierdo M, Elnagar R, Afonso J, Peñailillo L, et al. Effects of therapies involving plyometric-jump training on physical fitness of youth with cerebral palsy: a systematic review with meta-analysis. *Sports.* 2024;12(6):152. <https://doi.org/10.3390/sports12060152>.
45. Wang R, Li Q, Xue W. Effects of small-sided game training on lower limb explosive strength in handball players: a single-arm meta-analysis. *Front Sport Act Living.* 2024;6:1477347. <https://doi.org/10.3389/fspor.2024.1477347>.
46. Huang D, Wang X, Gonjo T, Takagi H, Huang B, Huang W, et al. Effects of creatine supplementation on the performance, physiological response, and body composition among swimmers: a systematic review and meta-analysis of randomized controlled trials. *Sports Med Open.* 2024;10(1):115. <https://doi.org/10.1186/s40798-024-00784-8>.

47. Herbison P, Hay-Smith J, Gillespie WJ. Meta-analyses of small numbers of trials often agree with longer-term results. *J Clin Epidemiol.* 2011;64(2):145–53. <https://doi.org/10.1016/j.jclinepi.2010.02.017>.
48. Jackson D, Turner R. Power analysis for random-effects meta-analysis. *Res Synth Methods.* 2017;8(3):290–302. <https://doi.org/10.1002/jrsm.1240>.
49. Ramirez-Campillo R, Sortwell A, Moran J, Afonso J, Clemente FM, Lloyd RS, et al. Plyometric-jump training effects on physical fitness and sport-specific performance according to maturity: a systematic review with meta-analysis. *Sports Med Open.* 2023;9(1):23. <https://doi.org/10.1186/s40798-023-00568-6>.
50. Ramirez-campillo R, Garc-Hermoso A, Moran J, Chaabene H, Negra Y, Scanlan AT. The effects of plyometric jump training on physical fitness attributes in basketball players : a meta-analysis. *J Sport Heal Sci.* 2022;11:656–70. <https://doi.org/10.1016/j.jshs.2020.12.005>.
51. McKay AKA, Stellingwerff T, Smith ES, Martin DT, Mujika I, Goosey-Tolfrey VL, et al. Defining training and performance caliber: a participant classification framework. *Int J Sports Physiol Perform.* 2022;17(2):317–31. <https://doi.org/10.1123/ijspp.2021-0451>.
52. Nawed A, Khan IA, Jalwan J, Nuhmani S, Muaidi QI. Efficacy of FIFA 11 + training program on functional performance in amateur male soccer players. *J Back Musculoskeletal Rehabil.* 2018;31(5):867–70. <https://doi.org/10.3233/BMR-171034>.
53. Fofha BM, Schwesig R, Ltifi MA, Bartels T, Hermassi S, Aouadi R. A 10-week FIFA 11+ program improves the short-sprint and modified agility T-test performance in elite seven-a-side soccer players. *Front Physiol.* 2023;14:1236223. <https://doi.org/10.3389/fphys.2023.1236223>.
54. Asgari M, Alizadeh MH, Shahrbanian S, Nolte K, Jaitner T. Effects of the FIFA 11+ and a modified warmup programme on injury prevention and performance improvement among youth male football players. *PLoS ONE.* 2022;17(10 October):1–11. <https://doi.org/10.1371/journal.pone.0275545>.
55. Ayala F, Pomares-Noguera C, Robles-Palazón FJ, Del Pilar García-Vaquero M, Ruiz-Pérez I, Hernández-Sánchez S, et al. Training effects of the FIFA 11+ and Harmoknee on Several neuromuscular parameters of physical performance measures. *Int J Sports Med.* 2017;38(4):278–89. <https://doi.org/10.1055/s-0042-121260>.
56. Daneshjoo A, Mokhtar AH, Rahnama N, Yusof A. The effects of comprehensive warm-up programs on proprioception, static and dynamic balance on male soccer players. *PLoS One.* 2012;7(12):e51568. <https://doi.org/10.1371/journal.pone.0051568>.
57. Zarei M, Abbasi H, Daneshjoo A, Barghi TS, Rommers N, Faude O, et al. Long-term effects of the 11+ warm-up injury prevention programme on physical performance in adolescent male football players: a cluster-randomised controlled trial. *J Sports Sci.* 2018;36(21):2447–54. <https://doi.org/10.1080/02640414.2018.1462001>.
58. Robles-Palazón FJ, Pomares-Noguera C, Ayala F, Hernández-Sánchez S, Martínez-Romero MT, De Baranda SP, et al. Acute and chronic effects of the Fifa 11+ on several physical performance measures in adolescent football players. *Eur J Hum Mov.* 2016;36:116–36.
59. Costa Silva JRL, Silva JF, Salvador PCdoN, Freitas CDlaR. O efeito do “FIFA 11+” na performance de saltos verticais em atletas de futebol. *Rev Bras Cineantropom Desempenho Hum.* 2015;17(6):733. <https://doi.org/10.5007/1980-0037.2015v17n6p733>.
60. Gatterer H, Lorenzi D, Ruedl G, Burtcher M. The, “FIFA 11+” injury prevention program improves body stability in child (10 year old) soccer players. *Biol Sport.* 2018;35(2):155–8. <https://doi.org/10.5114/biolSport.2018.71604>.
61. Impellizzeri FM, Bizzini M, Dvorak J, Pellegrini B, Schena F, Junge A. Physiological and performance responses to the FIFA 11+ (part 2): a randomised controlled trial on the training effects. *J Sports Sci.* 2013;31(13):1491–502. <https://doi.org/10.1080/02640414.2013.802926>.
62. Zhou X, Luo A, Wang Y, Zhang Q, Zha Y, Wang S, et al. The effect of FIFA 11+ on the isometric strength and running ability of young soccer players. *Int J Environ Res Public Health.* 2022;19(20):13186. <https://doi.org/10.3390/ijerph192013186>.
63. Akbari H, Sahebozamani M, Daneshjoo A, Amiri-Khorasani M. Effect of the FIFA 11+ programme on vertical jump performance in elite male youth soccer players. *Monten J Sports Sci Med.* 2018;7(2):17–22. <https://doi.org/10.26773/mjssm.180903>.
64. Hwang J, Kim J. Effect of fifa 11+ training program on soccer-specific physical performance and functional movement in collegiate male soccer players: a randomized controlled trial. *Exerc Sci.* 2019;28(2):141–9. <https://doi.org/10.15857/ksep.2019.28.2.141>.
65. Rohmansyah NA, Hiruntrakul A. Do lipid profiles, body composition, and physical fitness improve after a FIFA 11+ training programme from obese women? *Sport Mont.* 2021;19(2):51–7. <https://doi.org/10.26773/smj.210609>.
66. Ramos AP, de Mesquita RS, Migliorini F, Maffulli N, Okubo R. FIFA 11+ KIDS in the prevention of soccer injuries in children: a systematic review. *J Orthop Surg Res.* 2024;19(1):413. <https://doi.org/10.1186/s13018-024-04876-9>.
67. Ejnisman B, Andreoli CV, de Castro Pochini A, Cohen M, Bizzini M, Dvorak J, et al. Shoulder injuries in soccer goalkeepers: review and development of a FIFA 11+ shoulder injury prevention program. *Open Access J Sports Med.* 2016;7:75–80. <https://doi.org/10.2147/OAJSM.S97917>.
68. Little T, Williams AG. Specificity of acceleration, maximum speed, and agility in professional soccer players. *J Strength Cond Res.* 2005;19(1):76–8. <https://doi.org/10.1519/14253.1>.
69. Sheppard JM, Young WB. Agility literature review: classifications, training and testing. *J Sports Sci.* 2006;24(9):919–32. <https://doi.org/10.1080/02640410500457109>.
70. Zen MZ, Setijono H, Kusnanik NW. The effect of single leg and double leg bounding exercise programme (BEP) on increasing physical abilities, strength, and agility. *Br Int Humanit Soc Sci J.* 2021;3(3):459–67. <https://doi.org/10.33258/biohs.v3i3.525>.
71. Yalfani A, Saki F, Kerman MT. The effects of the FIFA 11+ and 11+ kids training on injury prevention in preadolescent football players: a systematic review. *Ann Appl Sport Sci.* 2020;8(4):2–8. <https://doi.org/10.29252/aassjournal.832>.
72. Kim H, Lee J, Kim J. The impact of the FIFA 11+ program on the injury in soccer players: a systematic review. *Asian J Kinesiol.* 2020;22(4):55–61. <https://doi.org/10.15758/ajk.2020.22.4.55>.
73. Bizzini M, Dvorak J. FIFA 11+: an effective programme to prevent football injuries in various player groups worldwide—a narrative review. *Br J Sports Med.* 2015;49(9):577–9. <https://doi.org/10.1136/bjsports-2015-094765>.
74. McKay CD, Steffen K, Romiti M, Finch CF, Emery CA. The effect of coach and player injury knowledge, attitudes and beliefs on adherence to the FIFA 11+ programme in female youth soccer. *Br J Sports Med.* 2014;48(17):1281–6. <https://doi.org/10.1136/bjspo-2014-093543>.
75. Nguyen KP, Zhou W, McKenna E, Colucci-Chang K, Bray LCJ, Hosseini EA, et al. The 24-h savings of adaptation to novel movement dynamics initially reflects the recall of previous performance. *J Neurophysiol.* 2019;122(3):933–46. <https://doi.org/10.1152/jn.00569.2018>.

76. Dello Iacono A, Beato M, Unnithan V. Comparative effects of game profile-based training and small-sided games on physical performance of elite young soccer players. *J Strength Cond Res*. 2021;35(10):2810–7. <https://doi.org/10.1519/JSC.0000000000003225>.
77. Haugen T, Seiler S, Sandbakk Ø, Tønnessen E. The training and development of elite sprint performance: an integration of scientific and best practice literature. *Sports Med Open*. 2019;5(1):44. <https://doi.org/10.1186/s40798-019-0221-0>.
78. Romero C, Ramirez-Campillo R, Alvarez C, Moran J, Slimani M, Gonzalez J, et al. Effects of maturation on physical fitness adaptations to plyometric jump training in youth females. *J Strength Cond Res*. 2021;35(10):2870–7. <https://doi.org/10.1519/JSC.0000000000003247>.
79. Bartolomei S, Grillone G, Di Michele R, Cortesi M. A comparison between male and female athletes in relative strength and power performances. *J Funct Morphol Kinesiol*. 2021;6(1):17. <https://doi.org/10.3390/jfmk6010017>.
80. Sugimoto D, Myer GD, Bush HM, Klugman MF, Jennifer MMM, Hewett TE. Compliance with neuromuscular training and anterior cruciate ligament injury risk reduction in female athletes: a meta-analysis. *J Athl Train*. 2012;47(6):714–23. <https://doi.org/10.4085/1062-6050-47.6.10>.
81. Sugimoto D, Mattacola CG, Bush HM, Thomas SM, Barber Foss KD, Myer GD, et al. Preventive neuromuscular training for young female athletes: comparison of coach and athlete compliance rates. *J Athl Train*. 2017;52(1):58–64. <https://doi.org/10.4085/1062-6050-51.12.20>.
82. Cormie P, McGuigan MR, Newton RU. Developing maximal neuromuscular power: Part 2 training considerations for improving maximal power production. *Sport Med*. 2011;41(2):125–46. <https://doi.org/10.2165/11538500-000000000-00000>.
83. Thomas K, French D, Hayes PR. The effect of two plyometric training techniques on muscular power and agility in youth soccer players. *J Strength Cond Res*. 2009;23(1):332–5. <https://doi.org/10.1519/JSC.0b013e318183a01a>.